

Silicon as a Stress Alleviator in Horticultural Crops: A Review

¹Rashid Hussain, ¹Muhammad Rashid Shaheen, ²Sahar Rashid, ¹Muhammad Abdul Rahman Iqbal, ³Qasim Ali, ⁴Muhammad Imran Ghani

¹Department of Horticultural Sciences, Faculty of Agriculture and Environment, The Islamia University of Bahawalpur, Bahawalpur 63100, Punjab, Pakistan

²Horticultural Research Institute, Ayub Agricultural Research Institute, Faisalabad, Punjab, Pakistan

³Department of Soil Science, Faculty of Agriculture and Environment, The Islamia University of Bahawalpur, Bahawalpur 63100, Punjab, Pakistan

⁴College of Life Sciences, Guizhou University, Guiyang 550005, China

About the Article

Open Access

Review Article

How to Cite: Hussain R, Shaheen MR, Rashid S, Iqbal MAR, Ali Q, Ghani MI. Silicon as a stress alleviator in horticultural crops: A Review. *Acta Agric. Sci.* 2025;1:1–19.

Keywords:

Abiotic stresses, phytohormones, silicon, stress tolerance

Corresponding author:

Rashid Hussain, Department of Horticultural Sciences, Faculty of Agriculture and Environment, The Islamia University of Bahawalpur, Bahawalpur 63100, Punjab, Pakistan

ABSTRACT

The persistence of agricultural systems is facing more complex threats due to environmental discontinuities, which makes the case for targeted approaches to ameliorate the resilience of crops to non-living environmental stressors. Among current practices, which range from mineral addition to conventional farming, silicon (Si) is viewed as an element of lesser importance to the agricultural industry, yet fundamental with regards to explains the adaptation of plants to the environment. The concentration of Si elements in most plants is low but Si elements are usually present, by adding it externally, adverse environmental conditions will easily increase, leading to decreased (osmotically) regulated conditions, stabilizing of cell membranes and oxidative stress. In contrast to other supplements, Si has the unique property of regulating metabolism of phytohormone. Phytohormones, including auxins, cytokinins, ethylene, gibberellins, salicylic acid, abscisic acid, brassinosteroids and jasmonate, are responsible for the communication and control of plant activity in response to external and internal environmental and developmental changes. As Si exists in plant systems, it increases hormone synthesis and receptor activity, antagonize hormone turnover and receptor activity, thereby organizing rationally coordinated defense responses to stress. Smart hormonal changes, in turn, has been positively associated with enhanced physiological integrity and defense responses. Silicon-driven stress resilience versatility includes additional multidimensional remodeling systems, including organizational, transcriptional, biochemical and even metabolic, beyond the aforementioned primary signaling. In Si-treated plants, such systemic stress tolerance modules impact the emerging integrated gene-protein-metabolite networks outlined by multi-omics systems frameworks. In terms of sustainable intensification, the Si integrated hormonal order is a realistic approach to food security and resilient horticultural production. In addition to interfaces of sustainable development, Si provides a new spine to internal control that mitigates abiotic stress. Future work is needed to demarcate threshold application levels, the timing of application and specific pathways toward individual crops. Translational silicates frameworks that draw on delineated system interfaces with advanced biotechnologies offer a critical avenue for the production of engineered stress-resilient crops to sustain agricultural yields amidst growing climate extremes.

INTRODUCTION

For survival and health promotion, it is imperative to ensure that nutritious and secure meals are available. However, the agriculture industry may encounter significant hurdles in the future in satisfying the food production demands of a rapidly growing human population. Over the next 30 years, the global population is predicted to increase by 2 billion people, reaching 9.8 billion by 2050¹. This population growth poses significant limitations on land availability, particularly in Asian countries². Additionally, human activities, including the release of greenhouse

© The Authors, 2025. Published by the Academia Publications. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (CC-BY) 4.0. (<http://creativecommons.org/licenses/by/4.0/>)

gases, contribute to the warming of the environment. This phenomenon adversely affects food security by disrupting resource availability and livelihoods of individuals³. As a result, the agriculture industry and scientists are constantly challenged to feed millions of people⁴. Abiotic stressors caused or exacerbated by climatic conditions account for nearly 50% of agricultural losses, making sustainable agricultural output vital to guaranteeing food security⁵. Plant growth rate and output are significantly influenced by temperature, humidity, light, moisture, salt, availability of nutrients and the concentration of industrial and agricultural chemicals in water and soil resources. Osmotic stress is caused by abiotic circumstances, which disturb ion transport and cellular equilibrium in plants. Furthermore, the interplay of abiotic and biotic stressors renders plants more vulnerable to pests⁶.

The plant root is home to a rich microbial community that forms complicated relationships with plants, impacting plant growth and production through metabolic characteristics and interactions^{7,8}. Plant-associated bacteria display morphological changes within the rhizosphere, reflecting a variety of groups suited to environmental circumstances, aiding adaptation to environmental challenges and maintaining plant health and resilience⁹. Growth-promoting rhizobacteria of plants (PGPRs) are soil bacteria that colonise plant roots (rhizosphere) and contribute to higher plant performance and yields¹⁰. The PGPRs not only form positive connections with roots, resulting in considerable growth responses but they also play an important function in phytopathogen regulation. They also help to improve soil structure and crop productivity by stimulating the development of soil aggregates and pores¹¹. Silicon (Si) use in agriculture has the potential to reduce abiotic stressors in crops¹². Silicon is the element with the eighth greatest abundance in nature and, after molecular oxygen, the second most widespread constituent in soil¹³. Si is found in soil as monosilicic acid (H_4SiO_4), which is uncharged at pH 9¹⁴. It may ionize into silicate ions, such as $(OH)_3 SiO_2$, at higher pH values (>9). Weathering of silicate-containing minerals releases Si into the soil¹⁵. Because most soils are naturally rich in Si, the availability of dissolved Si varies with soil type and conditions in the environment. However, soil types and seasonal fluctuations impact plant absorption. Silicon rapidly interacts with oxygen (O_2) to generate silicon dioxide (SiO_2), which accounts for around 50% of soil mass¹⁶. Si is found mostly in mineral soils such as SiO_2 and either primary or secondary silicate minerals, which also contain elements like Al (aluminosilicates) and Mg¹⁷. Recognizing Si's helpful effect in reducing abiotic stress in crops necessitates further research into its physiological and molecular features. This chapter seeks to offer a thorough grasp of Si's vital function in promoting plant growth and production in the face of

abiotic stress. Several studies have also been conducted to investigate Si's function in phytohormone synthesis, gene expression regulation and the impacts of environmental stress on metabolism and development. By exploring Si's potential, this chapter hopes to motivate additional study and reveal its critical role in ensuring global food security.

The importance of silicon for sustainable agricultural growth: The growing population has put enormous strain on the agricultural sector to solve food insecurity. While technological interventions have enhanced crop yields, industrialization has also resulted in increasing environmental pressures, which have harmed crop output and hampered long-term agricultural progress. Plants respond by modifying their metabolic networks and producing physiological changes¹⁸. Numerous studies have shown that Si has an important function in improving agricultural productivity and yield under adverse environmental circumstances¹⁹. Si buildup in plant organs improves both monocot and dicot crop growth and metabolism²⁰. Because of increased Na^+ competition with K^+ and Ca^{2+} transport in cell plasma membranes, for example, affects reproductive activities²⁰. According to recent studies, Si reduces Na^+ buildup, Osmotic pressure and oxidative damage^{21,22}. To counteract abiotic stressors, Si stimulates defense pathways linked with phytohormone production and compatible solutes^{23,24}. Exogenous Si treatment improves salt tolerance in salt-sensitive okra genotypes by increasing proline accumulation²⁵. Si also increases the synthesis of the total number of free amino acids, proline and glycine betaine, for example, are found in *Capsicum annuum* cultivars, improving salt stress resistance²⁶. Furthermore, Si buildup in crops such as tomato, okra, rice and wheat leads to a decrease in proline concentration, suggesting Si's efficiency in activating its own salt-resistance mechanisms²⁷. Silicon (Si) treatment improves crop growth, production and fruit quality in a variety of fruit crops^{28,29}. Si promotes water and energy conservation, improves fruit quality and boosts production. On the other hand, plants accumulate Si in distinct ways, resulting in diverse Si-mediated responses to abiotic stresses³⁰. Major crop plants, such as cotton, soybean, cucurbits and tomato as well as monocots like rice, maize and wheat, collect Si via specialized transporters to ameliorate diverse stress effects¹².

Si treatment increases the expression and activity of defense-related genes while also boosting the accumulation of defensive chemicals such as phenol compounds, phytoalexins and momi-lactones³¹. Supplementation of Si, for example, lowers Reactive Oxygen species (ROS) scavenging capabilities in vegetable cultivars such as *Cucumis sativus* under stressful circumstances³². Exogenous Si supplementation alleviates sulphur and osmotic stress in tomatoes by increasing root sucrose levels³³. Si can also

alleviate heavy metal stress by activating several pathways and scavenging ROS³³. Si enrichment increases plant growth and development by increasing fruit Nutrient Usage Efficiency (NUE)³⁴.

Silicon (Si) further mitigates the toxicity of metal ions in plants by either reducing their solubility or catalytically detoxifying them in the apoplastic space. Evidence indicates that Si enhances the tolerance of crops to UV-B stress by subsequent stimulation of defensive metabolites and mechanical thickening of the cell wall³⁵. Collectively, the data imply that Si exerts a constitutive and adaptive influence over the physiological and metabolic responses of crops exposed to prolonged water deficits. The preceding analysis focused on the modulatory role of Si in optimizing phytohormonal signaling pathways that regulate abiotic stress. Taken in aggregate, these observations underscore Si's pivotal contribution to elevating plant tolerance to chronic abiotic shock, thereby constituting a promising agronomic intervention to sustain productivity in environments afflicted by recurring, pronounced drought.

Differential synthesis of phytohormones in response to silicon and abiotic stress: The plant hormones Gibberellins, Cytokinins, Ethylene, Brassinosteroids, Jasmonic acid, Salicylic acid and Abscisic acid, along with signaling molecules such as Calcium and Nitric oxide, exert pronounced influence on physiological processes, even when present in minor concentrations³⁶. Within horticultural science, these phytohormones have been implicated in the growth and morphogenesis of economically important vegetable crops, including tomato (*Solanum lycopersicum*), cucumber (*Cucumis sativus*) and pepper (*Capsicum annuum*)³⁷. Concurrently, silicon exhibits decisive interaction with reactive oxygen species to restore and preserve cellular symmetry at the structural, molecular and biochemical scales, thereby mitigating a wide spectrum of abiotic and biotic stresses. By modulating multiple pathways-nutrient assimilation, the regulation of source and sink dynamics and the reinforcement of cellular antioxidant and signaling networks-silicon measurably enhances the plant's resilience to severe environmental perturbations³⁸. Furthermore, combining critical metabolites with silicon to ameliorate abiotic stressors has shown complex and interrelated mechanisms that can improve crop development and production, as seen in pepper (*Capsicum annuum*)³⁹. The complex link between silicon-mediated plant growth and stress tolerance, which involves cross-talk with phytohormones, ROS and signaling molecules, has received much attention. Gibberellins (GA) are phytohormones that contribute considerably to many developmental phases in plants and act as effective mechanisms to counteract abiotic stressors⁴⁰. Higher dosages of silicon supplementation resulted in increased endogenous levels of GA₁ and GA₄ in

Crocus sativus plants subjected to salt and drought stress, showing a major metabolic reshuffling inside stressed plants following silicon therapy⁴¹.

Si treatment has been reported to promote the production and accumulation of Gibberellins in salt-stressed *Glycine max* and *Oryza sativa* plants while lowering the levels of Jasmonic Acid (JA) and Ethylene (ET)⁴². Apart from its importance in a variety of plant processes involved in development and growth, cytokinin also shows a significant role in effectively reducing abiotic stressors. Si-mediated intervention increases cytokinin production, which improves their stress-resistance function⁴¹. Cadmium (Cd) poisoning harms pea plants, producing changes in root and stem development, architecture and biochemistry. Plants exposed to Cd toxicity develop more slowly with Cd buildup, decreased chlorophyll content and changes in primary metabolites such as carbohydrates and proteins. Cadmium exposure induces oxidative stress in plant systems, evidenced by elevated malondialdehyde (MDA) and hydrogen peroxide (H₂O₂) levels, subsequently triggering an up-regulation of antioxidant enzymes, specifically catalase and peroxidase, in an adaptive response to cellular perturbation. Structural alterations noted under increasing cadmium concentrations include the development of aberrant vascular patterns, irregular lignification in the pith parenchyma and hypertrophy of cortical parenchyma, corroborating the morphological instability observed by El-Okkiah et al.⁴³. Conversely, pre-application of silicon ions effectively abrogates the oxidative burst associated with cadmium stress, thereby sustaining cellular homeostasis. The concomitant use of silicon and 24-epibrassinolide has been demonstrated to bolster *Brassica juncea*'s resilience to sodium chloride lethality by sustaining maximal photochemical efficiency, enhancing the activities of catalase, peroxidase and superoxide dismutase and by promoting the accumulation of proline, as reported by Siddiqui et al.⁴⁴. Moreover, silicon treatment positively modulates membrane integrity, an effect partly ascribed to an augmented biosynthesis of jasmonic acid, a lipid-derived phytohormone consistently elevated in stressed tissues, thereby reinforcing osmotic adjustment and stress signaling pathways, as elaborated by Bhardwaj et al.²⁴.

Categorization of physiological responses indicates that exogenous silicon mitigates saline stress by conferring cellular protective modalities. Such treatment preferentially lowers ion efflux and membrane lipid peroxidation, while concurrently depressing Jasmonic Acid (JA) concentrations relative to untreated cohorts; by contrast, levels of Indole-3-acetic Acid (IAA) and Salicylic Acid (SA) exhibit elevation⁴⁵. Profiling of defense signaling reveals that silicon augments transcripts participating in JA biosynthetic cascades in mechanically wounded tissues, thereby attenuating injury-induced stress. Nonetheless, a more

granular elucidation linking silicon-mediated modulation of JA-associated mRNA abundance to stress relief remains to be consummated⁴⁶. Concurrent work shows that conjoint application of Si and external salicylic acid confers acid stress tolerance to tomato seedlings by elevating endogenous sodium and potassium reservoirs; this osmotic adjustment in turn curtails Reactive Oxygen Species (ROS) accumulation through reinforcement of antioxidant pathways and by transcriptionally up-regulating transporters and enzymes of Si acquisition and SA biosynthesis⁴⁶. Salicylic Acid (SA), a phenolic phytohormone, is essential for dealing with abiotic stressors and sustaining optimum plant development⁴⁷. SA promotes the antioxidative defense mechanism in plants to reduce heavy metal toxicity, although its levels have decreased when Si is used on the surface⁴⁸.

Abscisic Acid (ABA), also known as the stress hormone, accumulates in plants subjected to salt stress, causing stomatal closure and consequent water conservation⁴⁹. ABA modulates a wide range of physiological reactions in vegetable plants, involving stomatal closure and seed dormancy, assisting in adaptation to abiotic conditions such as drought or excessive salt⁵⁰. For example, ABA stimulates stomatal closure in tomato plants (*Solanum lycopersicum*) under water stress⁵¹ and modulates seed dormancy in lettuce plants (*Lactuca sativa*) under unfavorable environmental circumstances⁵². An increased supply of Si to salt-stressed vegetable plants results in a beneficial relationship between Si activity and ABA. Exogenous Si supplementation improves salt tolerance by increasing Si absorption and ABA synthesis⁵³. In Si-mediated stress tolerance, ABA has an antagonistic connection with SA/JA biosynthesis, demonstrating ABA's dynamic involvement in response to Si treatment. Plant ABA content reduces when Si concentration increases during heavy metal, salt and drought stress conditions⁵⁴. Hormones that cause stress, such as ABA, JA and SA, are testified to rise in cold-stressed maize (*Zea mays*) seedlings, whereas the production of other plant growth regulators, such as auxins, gibberellins (GA) and cytokinins, is inhibited⁵⁵.

Polyamines, a group of nitrogen-rich, aliphatic cationic molecules that includes putrescine, spermidine and spermine, are known to accumulate in elevated amounts within plant tissues as a cellular strategy to mitigate oxidative and salinity stress⁵⁶. Empirical evidence supporting their protective role continues to mount, notably evidenced by enhanced synthesis when plants are treated with exogenous silicon (Si), a response thoroughly documented in *S. bicolor*⁵⁷. Following Si supplementation, transcriptional data show that the S-adenosyl-L-methionine decarboxylase (SAMDC) gene-coding for the rate-limiting enzyme in polyamine biosynthesis-exhibits elevated expression. The resultant increase in polyamine biosynthetic activity, driven by Si-mediated signal transduction, in turn

orchestrates a cascade of cytoprotective and metabolic buffering effects that collectively attenuate stress injury and promote the sustainment of vegetative and reproductive processes³¹.

Silicon's mechanism for reducing abiotic stress: Silicon exerts its influence on plants by modulating multiple developmental processes, thereby serving as a crucial element in mitigating diverse abiotic stresses. One of its primary mechanisms involves promoting the upregulation of antioxidants⁵³.

Silicon's role in enhancing salt tolerance in plants: Salinity stress has a detrimental influence on the development and visual appeal of ornamental plants and it continues to be a key limiting factor for agricultural output globally^{58,59}. Morphological features such as leaf number and leaf area can be harmed by salt stress⁶⁰. Furthermore, lower chlorophyll levels, relative amount of water, important nutrient intake and higher leakage of electrolytes are seen⁶¹. Silicon can boost production in salt-stressed plants⁶². Although silicon is not a required nutrient, it does engage in processes of metabolism that increase resistance in plants to environmental circumstances such as drought and salt⁶³⁻⁶⁵. Silicon application is a low-impact method for improving plant reactions to salt stress⁶⁶. It improves plant resistance to salinity and drought stress^{26,67}, minimizes the detrimental effects of salt stress on chlorophyll content and the generation of biomass⁶⁸ and promotes adaptive responses like phenolic compound manufacture, mineral uptake and antioxidant activity^{69,70}.

Under both salt stress and normal circumstances, halo-priming with silicon nanoparticles (Si NPs) encourages the development of seedlings and germination^{64,71}. Silicon nanoparticles (Si NPs) have received a lot of interest in agriculture, horticulture and biotechnology because of their ability to withstand environmental challenges and post-harvest losses⁷². Nanotechnology has had a revolutionary influence on horticulture, notably in fruit crops, due to its efficacy against environmental challenges and post-harvest losses⁷³.

The incorporation of nano-silica (nSiO₂) has been demonstrated to mitigate salt-induced plant damage while simultaneously facilitating faster vegetative growth in strawberry cultivars⁶². Halo-priming applications employing silicon or silicon nanoparticles (Si NPs) conferred enhanced salt tolerance in *Lathyrus* seedlings, according to findings by El-Serafy et al⁵⁸. In sweet pepper, an aerial silicon application significantly elevated the concentrations of both chlorophyll a and b, ameliorated the translocation of mineral nutrients, improved the hydration of the leaf tissue and ultimately increased fruit yield. Concurrent physiological advantages included decreased lipid peroxidation, reduced electrolyte leakage and lowered levels of superoxide and

hydrogen peroxide⁷⁴. In a wider context, silicon supplementation has been correlated with improved germination rates, faster germination percentages and lengthier shoot growth-benefits that registered in both unstressed and saline or drought-stressed scenarios^{75,76}. By contrast, the influence of silicon on saline-stressed root systems has yielded mixed or negative results. At the same time, shoot growth recovery has been recorded in wheat and cucumber, the root systems in those same trays exhibited negligible enhancement⁷⁷. In a complementary study of tomato, the employment of SiO_2 nanoparticles elicited appreciable physiological improvements, signifying a source of consistent positive responses from this class of novel silica amendments⁷⁸. Tantawy et al.⁷⁹ discovered that nano silicon is more effective and efficient at mitigating salt stress in sweet pepper plants. Spraying Si NPs on "Ewais" mango plants increased overall yield (number and weight of fruits per tree) and fruit physiochemical properties⁸⁰. To counteract salt stress in watermelon agriculture, a combination of mycorrhiza and silicon might be used⁸¹.

Si uses various methods to protect plants against salt stress. These strategies include minimizing ion toxicity while maintaining plant water balance, improving inorganic absorption and incorporation, controlling the production of suitable solutes and phytohormones, mitigating oxidative stress, changing gas exchange characteristics and manipulating gene expression. The reduction in Na^+ uptake and accumulation, which has been widely observed with Si supply under salt stress, results in a boosted K^+/Na^+ ratio¹². However, the molecular basis of Si absorption in fruit trees is unclear⁸².

Silicon amendment mitigates oxidative damage in a variety of plant species facing saline stress by augmenting the activity of key antioxidant enzymes, including superoxide dismutase (SOD), catalase (CAT), peroxidase (POD), glutathione S-transferase (GSH) and ascorbate peroxidase (APX), while stabilizing membrane fluidity and sustaining plasma membrane H⁺-ATPase function, all of which culminates in the suppression of Reactive Oxygen Species (ROS) accumulation^{83,84}. In addition to preventing oxidative injury, Si application enhances salt tolerance by reprogramming osmotic adjustment mechanisms in the shoots and roots of okra, as evidenced by modifications in the concentrations of proline, glycine betaine and the pool of free amino acids²⁵. Salt-stressed soybean seeds, in turn, exhibit altered phytohormonal profiles, notably a rise in gibberellin (GA) titers concomitant with reduced abscisic acid (ABA) levels, following Si supplementation⁸⁵. The fabrication and administration of nano-silicon dioxide in strawberry plants positively impacted the stability of the epicuticular wax layer, preserved chlorophyll and carotenoid concentrations and restrained proline accumulation⁶². Such effects, in concert, enabled the amelioration of morphological and metabolic derangements customary to

saline environments⁶¹. Furthermore, the accumulation of silica in plant tissues has been observed to correlate with the surveillance of epicuticular wax biosynthesis; for example, the exposure to Si treatments modified the morphology of fruit trichomes in cucumber (*Cucumis sativus* L., cv. Corona) such that a significant fraction of silica deposit was sequestered within the wax of trichome cells⁸⁶.

Silicon's contribution to drought resistance in plants:

Drought is a key environmental restriction that limits the development and production of field and horticultural crops^{87,88}. It is distinguished by limited water availability, both physically and physiologically, which has a negative worldwide influence on the evolution of plants and output⁸⁹. Drought-induced stressors have a detrimental influence on agricultural yield and quality⁹⁰, particularly in arid and semiarid locations where water shortage stress is a major factor impacting crop productivity⁹¹. Water deficit stress response varies depending on plant species, development stage, interval and strength of the stress⁹². Globally, significant efforts have been undertaken to increase irrigation water efficiency in diverse fruit crop species⁹³.

Drought-induced osmotic stress, on the other hand, increase the formation of Reactive Oxygen Species (ROS)⁹⁴, which can cause oxidative damage to proteins, DNA and lipids⁹⁵. Prolonged water shortages can impair stomatal sensitivity to low water potential, resulting in lower water content, turgor and total water potential, as well as partial or full stomatal closure. Cell enlargement, growth rate and total plant growth are all lowered under these circumstances⁹⁶. Plants contain both enzymatic and non-enzymatic antioxidant systems that help them endure stress caused by water and avoid photooxidative harm⁹⁷.

Agriculture researchers have long explored strategies to reduce the detrimental effects of drought on crops. Exogenous protectants, which include trace elements such as silicon (Si), are a regularly used approach. Plants that store silicon can endure water stress by using efficient compounds. Silicon has been shown to be beneficial in reducing biotic pressures such as plant disease and insect pests, drought, salt, metal toxicity, temperature stress, nutritional imbalance and waterlogging are among abiotic difficulties. Its significance has been recognized in the development of horticultural crops such as vegetables, fruits and floricultural crops. Plants treated with silicon have higher stomatal conductance, transpiration rate, Leaf Relative Water Content (LRWC) and hydraulic conductivity in the roots and throughout the plant. When used in areas with limited water supply, silicon increases LRWC, improves water potential through osmotic adjustment, decreases electrolyte leakage and impacts proline levels. Furthermore, silicon boosts the antioxidant defense system in drought-stressed plants. The key mechanism behind silicon's protective role in plants is the buildup of polycyclic acids in cells⁹⁸.

Research demonstrates that the promotion of silicon in water-stressed crops and cultivars can exhibit cultivar-specific and cultivation-condition labile morphological and physiological alterations⁹⁹; for instance, Si supplementation has been associated with enhanced radicular architecture in melon¹⁰⁰. Such effects extend beyond structural adaptation: the element intervenes in a constellation of biochemical, physiological and photosynthetic pathways, synergistically fortifying the plant against the rigours of hydric deficit¹⁰¹. During episodes of water shortage, Si has been shown to sustain germination rates, permit greater biomass accumulation and elevate the photosynthetic carbon-assimilation rate, mechanisms that are correlated with osmotic adjustment, modulation of stomatal conductance, improved mineral translocation and a robust redox balance governed by modulated reactive oxygen species scavenging⁶⁷. Notwithstanding its earth-abundant and insoluble bulk, silicon, constituting circa 32% of the crust's mass, enters the plant through the mineralised reaction mediated by the mono-silicic acid monomer^{102,103}.

Accumulating evidence supports a beneficial role for silicon as an exogenous stress-relief agent across diverse horticultural species, encompassing tomatoes (*Solanum lycopersicum* L.), carrots (*Daucus carota* L.) and melon¹⁰⁴⁻¹⁰⁷. When applied as a soil or foliar amendment, silicon markedly alleviates imposed drought stress, as observed in cantaloupe (*Cucumis melo* L.) and strawberry (*Fragaria ananassa* Duch.)¹⁰⁸. Notably, the Se/SiO₂ composite has been highlighted as an effective formulation in field trials, ameliorating soil moisture deficits in strawberry and additional agronomic species¹⁰⁹.

Subsequent phytochemical analyses reveal that, in response to drought stress, the foliar tissues of all tested mango cultivars exhibit a consistent enhancement in the activity of key antioxidative enzymes, specifically peroxidase (POX), catalase (CAT) and superoxide dismutase (SOD), across a full three-season trial¹¹⁰. Silicon has an effective mechanism for scavenging reactive oxygen species, which protects plants from oxidative damage and improves their capacity to endure environmental stress in dry environments, including mango trees¹¹⁰. The administration of Na₂SiO₃ has been demonstrated to be an effective therapy for severe water shortage stress, notably in the '*Gavioita*' strawberry cultivar¹¹¹. However, silicon inhibited the generation of activated oxygen species and oxidative damage in potato plants subjected to water stress¹¹². Water scarcity is a severe abiotic stress that can reduce tomato output by up to 50%¹¹³. Exogenous silicon treatment reduces the antagonistic effects of drought stress in tomatoes¹¹⁴ and greatly enhances tomato crop water usage efficiency¹¹⁵. Cantaloupe production, like other horticultural crops, needs adequate management practices to obtain a good yield and high-quality fruit¹¹⁶. Low-quality fruit

development in cantaloupes is frequently attributed to insufficient pollination, prevailing high temperatures and water-deficit stress conditions¹¹⁷. Application of Si-based fertilizers, when applied at appropriate rates in conjunction with regulated soil moisture, has been shown to enhance yield and irrigation water productivity, thus suggesting agronomic advantages for cantaloupe cultivation when subjected to mild drought stress¹⁰⁸. Further corroborative evidence indicates that exogenous applications of magnesium carbonate, enriched with silicon, in banana cultivation can function effectively as an anti-transpirant agent, thereby augmenting fruit weight and improving overall fruit quality¹¹⁷.

Silicon's influence on heavy metal stress mitigation: Toxic metalloids such as copper, lead, nickel, chromium, mercury, zinc, arsenic and cadmium are contaminating a substantial agricultural land area due to urban or peri-urban cities¹¹⁸. Pollution from the environment or human activity can both introduce heavy metals into the food chain¹¹⁹. Heavy metal contamination of agricultural land can occur because of the use of sewage, sludge or industrial effluents¹²⁰. Consuming a variety of fruits and vegetables is critical for receiving key micronutrients and dietary fibre for overall wellness. According to American standards for 2015-2020, fruits and vegetables should account for around half of every single meal¹²¹. In research, toxic metal exposure in fruits and vegetables has been documented due to fertilization, irrigation and air deposition¹²². Consuming contaminated veggies can expose you to heavy metals and offer a serious health risk¹²³. Heavy metals in the soil can be taken up by plants and delivered to pollinators during pollination, thereby changing their behaviour and plant fitness¹²⁴. Heavy metal contamination has also harmed the quality and safety of medicinal plants¹²⁵. Consuming medicinal plants polluted with lead (Pb), mercury (Hg) and copper (Cu) are examples of heavy metals endangers consumer health. To solve this issue, it is critical to manage heavy metal levels in medicinal plants¹²⁵. Certain hazardous non-essential heavy metals, such as Pb, Hg, Cd and As, can be harmful to human health even at low levels, particularly in susceptible populations such as pregnant women and young children¹²⁵⁻¹²⁷. Increased Cd intake can harm the liver, respiration, kidneys and bones¹²⁸. Elevated Pb levels in the circulation can cause hypertension, compromised skeletal, immunological and endocrine systems, decreased cognitive ability in children and changes in renal and cardiac functions in adults¹³⁰. As a result, it is critical to analyse the human health concerns associated with heavy metals in widely consumed fruits and vegetables¹²⁸. In addition, extended exposure of grown crops to metal-contaminated growth medium reduces production sustainability. Toxic metal-induced soil fertility loss and nutrient imbalances lead to reduced agricultural yields^{130,131}.

Heavy metals including chromium, nickel, copper, arsenic, cadmium and lead were detected in potato (*Solanum tuberosum*), red onion (*Allium cepa*) and wild carrot (*Daucus carota*) in a study conducted by Islam et al.¹³². Heavy metal pollution reduced the chlorophyll content and biomass of vegetables, according to Li et al.¹³³. Peroxidase activity, on the other hand, rose at low concentrations and decreased at high concentrations of heavy metals. They advocated for the development of salt-tolerant crops such as tomato (*Solanum lycopersicum*) in heavy metal-contaminated regions. Green pepper (*Capsicum annuum*) and lettuce (*Lactuca sativa*) collected high levels of copper and zinc, whereas green pepper, tomato and onion acquired considerable quantities of lead¹³⁴. These heavy metal deposits were linked to the development of tannery enterprises in the region. Another research conducted by Lacatusu and Lacatusu¹³⁵ found significant levels of Cd and Pb in carrots (*Daucus carota subsp. sativus*), radish (*Raphanus raphanistrum subsp. sativus*) and potatoes (*Solanum tuberosum*). Cd and Pb levels in lettuce, parsley, dill and orach were up to 17 times higher than the allowable limits.

Silicon has been found to reduce the negative impacts of heavy metal pollution as well as long-term dangers in agriculture^{136,137}. Heavy metal removal by silicon may be divided into two categories: external and internal^{135,138}. External approaches include metalloid absorption or inactivation by silicon, as well as changing their present form by the addition of silicate minerals. Internal techniques, on the other hand, include processes such as increasing antioxidant enzyme activity, complex formation and compartmentalization, all of which assist lessen the negative consequences of heavy metal toxicity¹³⁸.

Heightened transcription of metal ion transporters such as OsLsi and OsHMA3 has been implicated in the mechanisms whereby Si amendment ameliorates Cu and Cd toxicity. Evidence further reinforces this proposition, with recent studies linking Si application to the induction of pathways that collectively scavenge reactive oxygen species¹³⁹. These pathways decrease the bioavailability of toxic metals either through the precipitative immobilization of cations within the cell wall matrix, or via biochelation and detoxification processes that effectively sequester metal ions and alter their cellular trafficking bioavailability^{140,141}. In peanut and rice seedlings, Si has been shown to lower the root, stem and foliar uptake of aluminum and chromium, further corroborating its protective action against the deleterious impacts of heavy metals¹⁴². Moreover, Barley exposed to osmotic and sulphur imbalances exhibited and enhanced root system biomass, accompanied by stratospheric sucrose concentrations consequent to Si foliar amendment, mitigating the detrimental stresses attributed to abnormal osmotic conditions and sulphur deficiency¹⁴³. The attenuation of metal ion activity by Si proceeds via

a combinatorial suppression of soil acidity, K, Ca, or Al fluctuation, selective precipitate formation and cellular reservoir routing¹⁴⁴. Quantitative expression assays demonstrate that exogenous Si significantly elevates PsbY transcription (Os08g02530), a polyprotein contributor to the antenna complex of Photosystem II, whilst zinc elicitation exhibits the converse expression profile in comparable rice conditions. Plants primarily absorb available soil Si as monosilicic acid through their roots^{145,146}.

Silicon is plentiful in practically all soil types, accounting for roughly 28% of soil composition¹⁴⁷. However, a large amount of silicon occurs as insoluble silicon dioxide, which is inaccessible to plants. Furthermore, some soils may be lacking in monosilicic acid, which is required by plants to satisfy their silicon requirements¹⁴⁸. Silicon fertiliser has been proven to boost total plant biomass by improving morphological and physiological parameters, as well as fibre quality¹²⁰. Furthermore, silicon fertilisation has been shown to reduce acid rain's negative effects on rice plants¹²⁰. The removal of heavy metals by silicon may be divided into two categories: internal and exterior. External approaches include metalloid absorption or inactivation, enhancing metal activity, or changing their present form by adding silicate elements. Internal techniques, on the other hand, include processes such as increasing antioxidant enzyme activity, complex formation and compartmentalization, all of which assist lessen the negative consequences of heavy metal toxicity¹³⁹.

Promoting wound stress tolerance with silicon: Environmental stress involves any external factor capable of perturbing living systems¹⁴⁹. The disruption of tissue integrity in cut fruits and vegetables accelerates both respiration and ethylene synthesis, outcomes intensified by the greater surface area-to-volume ratio and concomitant water efflux¹⁵⁰. Plant wound stress arises from abiotic and biotic agents, including mechanical lesions induced by wind or herbivory¹⁵¹. The compromised parenchyma, vasculature and epidermis become susceptible to secondary colonization by bacteria and fungi³¹. In both C3 and C4 crop species, lodging events represent a common wounding factor¹⁵². The molecular patterns elicited by mechanical injury, similar to salt stress, promote a heightened oxidative burst, damaging the polyunsaturated acyl moieties of the membrane lipids and resulting in membrane destabilization.

Upon mechanical injury, tissue-derived lipoxygenase catalysis escalates membrane lipid peroxidation, thereby intensifying Jasmonic Acid (JA) synthesis via α -linolenic acid-derived signals¹⁵³. Notably, mechanical wounding amplifies both lipid peroxidation and JA biosynthesis¹⁵⁴. A rigorous comprehension of JA-mediated pathways that underpin stress tolerance during wound events is, therefore, imperative. Several lines of inquiry have recently targeted silicon supplements, demonstrating lodging mitigation in

both rice and wheat^{155,156}. Site-specific disruption triggers oxidative bursts in affected cellular masses. Subsequent measurements indicate that catalase (CAT), peroxidase (PO) and polyphenol oxidase (PPO) activities in rice subjected to silicon amendment and mechanical injury exhibit significantly elevated levels relative to untreated controls¹⁴⁰. Corresponding increases in antioxidant enzyme activities have likewise been documented in wounded wheat¹⁵⁷, maize¹⁵⁸ and barley¹⁵⁸. The hydrogen peroxide (H_2O_2) generated is phytotoxic; however, silicon-acclimated tissues demonstrate elevated pools of CAT and PO that catalyse the conversion of H_2O_2 to innocuous water under wounding. Consequently, the augmented synthesis of antioxidative enzymes under silicon treatment serves as an effective defensive stratagem against the secondary oxidative stress that accompanies mechanical injury.

Research indicates that silicon applications mitigate lipid peroxidation during wounding events while simultaneously modulating oxidative stress¹⁵⁹. It is already established that plant cultivars exhibit variable physiological and metabolic responses to both biotic and abiotic stressors and such divergences extend to the composition and quantity of bioactive metabolites¹⁶⁰. Notably, Si-amended rice cultivars exhibit significantly reduced levels of lipid peroxidation markers following mechanical stress, in contrast to untreated controls¹⁴⁰. Elevated leaf silicon concentrations lead to a distinctive development of silicified cells, which exhibit a reduced degree of lipid saturation; this is corroborated by the noted decrease in malondialdehyde, a recognized end-product of lipid peroxidation¹⁶¹. The deposition of Si within cellular structures yields a hardened wall that serves as a mechanical barrier, thereby physically obstructing the ingress of potential pathogens through wound sites and thereby curtailing subsequent infection pathways¹⁶². In the context of interaction with herbivores, the high silicon concentration in the leaf tissue of rescue grass has been established to lead to substantial mandibular wear in grasshoppers, thus indicating an additional mechanical deterrent¹⁶³. Consequently, such findings collectively position silicon as a key mediator of physical resistance toward mechanical and subsequent biotic stress¹⁵⁹. Notably, the precise biochemical pathways that underlie the observed protective effects in both metabolic and structural contexts have yet to be elucidated, warranting further mechanistic investigation.

Enhancing high temperature stress tolerance with silicon: Anthropogenic climate change is raising heat stress, among the most severe abiotic pressures on terrestrial flora¹⁶⁴. Projections indicate mean global temperatures may increase by 1.5°C by the mid-century and 3.5°C by the close of the century¹⁶⁵. Phenological adjustments, including altered timing of budburst and leaf senescence, have been observed across numerous plant species subjected to rising

mean temperatures¹⁶⁶. Crop taxa sensitive to thermal extremes, such as *Prunus* and *Vitis*, as well as economically important overlaps of *Citrus* and various berries, remain particularly exposed¹⁶⁷. Elevated thermal regimes have been shown to perturb the maturation and colour development of *Citrus*, resulting in quality-limiting disorders, including insolation necrosis¹⁶⁸. Extreme thermal stress compromises cellular homeostasis by perturbing osmotic and ionic balances, thus inhibiting critical metabolic and physiological functions and by extension, curtailing biomass accumulation, developmental milestones and commercial viability¹⁶⁹. Episodes of excessive heat during anthesis and the establishment of the seed set stage have been associated with severe yield reductions, representing a major barrier to the improvement of global food security^{170,171}. Heat stress is known to depress vegetative growth, impair chlorophyll biosynthesis and diminish enzymatic activities vital to metabolic integrity¹⁷². Elevated leaf temperatures stimulate the excessive generation of Reactive Oxygen Species (ROS), resulting in oxidative cellular damage and the subsequent disruption of antioxidant and repair systems¹⁷³.

Improving horticultural plant microclimates is critical for guaranteeing year-round production¹⁷⁴. Foliar applications of silicon (Si) to mitigate heat stress within agricultural systems are widely adopted, with documented enhancements to yield quantity and quality¹⁷⁵. Bakhat et al.¹⁷⁵ endorse Si spraying as a sustained intervention for augmenting productivity in vegetable cropping systems. The physiochemical interactions triggered by Si deposition reduce heat-related metabolic perturbations, effectively buffering plants from exigent heat waves and as a consequence, safeguarding economic yield¹⁷⁶. In addition, Si behaves as a modulatory agent across critical physiological domains, influencing seed germination, stomatal dynamics, ionic translocation, membrane conductivity, photosynthetic efficiency and cumulative growth rates¹⁷⁷. Cucumber (*Cucumis sativus* L.), acknowledged as among the most extensively cultivated greenhouse vegetables, experiences protracted biotic and abiotic stress in controlled environments¹⁷⁸. Exceeding recommended mineral nutrient applications, the progressive sodification of root substrates and recurrent peak-season thermal accumulation constitute substantial yield-limiting factors, particularly in arid greenhouse systems¹⁷⁹. The employment of Si foliar applications has been corroborated as an effective amelioration, dampening both salinity and thermal stress symptoms in cucumber, thereby sustaining plant metabolism and productivity¹⁸⁰.

The combination application of Si and salicylic acid has beneficial benefits on cucumber plants in minimising the negative effects of high temperatures¹⁸¹. Under salt and heat stress circumstances, Si supplementation improves photosynthetic metrics and chlorophyll content in cucumber leaves¹⁸⁰. Under heat stress circumstances, the combined use

of rice straw as a soil cover and potassium silicate treatment improves soil fertility, nutrient absorption and crop output¹⁸². Unfavorable weather conditions, climate change and nutrient imbalances can all have a detrimental impact on tomato growth and output¹⁸³. High summer temperatures can stymie tomato growth and reduce output¹⁶⁵. Tomato seedlings, in contrast, may withstand severe temperature stress in greenhouses by receiving foliar applications of plant probiotics, sucrose and silicon¹⁸⁴. Shading has been proven to boost plant development and production by minimising the negative effects of high summer air temperatures on tomato plants¹⁸⁵. The foliar spray of potassium silicate has a considerable effect on celery yield during heat stress¹⁷⁷.

High temperatures during the blooming stage of monocotyledonous and dicotyledonous plants such as *Brassica napus*, *Solanum lycopersicum* and *Triticum aestivum* can diminish seed output¹⁷². Furthermore, spraying magnesium carbonate (silicon) as an anti-transpiration treatment has been proven to increase growth parameters, yield weight and fruit qualities in bananas¹¹⁷.

By reducing sunburn, calcium silicate has been found to improve the value and worth of Valencia orange fruits¹⁶⁸. Exogenous silicon (Si) administration improves the physiological growth of *Salvia splendens* at high temperatures by boosting the antioxidant system¹⁸⁶. The date palm is well-known for its capacity to survive adverse environmental conditions such as salt, drought and heat stress. Heat and drought, on the other hand, can have a detrimental influence on fruit yield, quality and a variety of physiological and metabolic processes in date palm seedlings^{187,188}. GA₃ and Si coupled have been found to boost plant development and rescue date palm growth under heat stress conditions¹⁸⁹.

As a cool-season food legume, garden pea is quite sensitive to temperature changes¹⁹⁰. Even a one-degree temperature increase can be termed heat stress for cool-season food legumes, with serious consequences for their development and metabolic functioning¹³⁴.

Understanding the impact of silicon on cold stress: Cold and freezing stress has been shown to depress crop yields by as much as 70%¹⁹¹, primarily owing to the simultaneous decline of temperature and irradiance¹⁹². In certain species, suboptimal chill conditions induce large-scale physiological cascades, manifesting as abnormal leaf expansion, stunted growth and extensive necrosis¹⁹³. These conditions likewise accelerate the generation of Reactive Oxygen Species (ROS), including superoxide radical (O₂·) and hydrogen peroxide (H₂O₂), the accumulation of which evolves as a secondary stress¹⁹². Elevated ROS concentrations compromise cellular integrity by peroxidising membrane lipids and inducing programmed cell death¹⁹⁴. In mitigation, plants synthesize an array of enzymatic antioxidants,

including superoxide dismutase (SOD), catalase (CAT), guaiacol peroxidase (GPX), glutathione peroxidase (GSH-Px), ascorbate peroxidase (APX), glutathione reductase (GR), dehydroascorbate reductase (DHAR) and monodehydroascorbate reductase (MDHAR), all of which operate in an integrated defense network against ROS¹⁹⁵. Non-enzymatic antioxidants, namely reduced glutathione (GSH) and ascorbate (AsA), further reinforce this defense array, collectively maintaining the cellular redox homeostasis requisite for survival in chilling environments. Plant tolerance to cold stress is increased by supplementing with silicon (Si), which maintains photochemical processes, photosynthetic gas exchange and activates antioxidant defence capability¹⁹⁶⁻¹⁹⁸. Exogenous Si treatment raises silicon concentrations in plants¹⁹⁹ and increased endogenous silicon deposition lowers lipid peroxidation²⁰⁰. The quantity of malondialdehyde (MDA), which represents membrane damage, can be used to detect lipid peroxidation²⁰¹. Low temperature, for example, can increase MDA content^{32,202}, indicating membrane degradation under cooling temperatures and low light circumstances¹⁹². Silicon supplementation, applied via seed soaking, foliar spray, or root drench, alleviates chilling injuries and frost incidence in species regardless of their Si-accumulators capacity²⁰³. Its protective capacity extends across numerous abiotic stressors, including lodging, drought, excessive radiation, thermal extremes and biochemical stressors such as salinity, metallic toxicity and nutrient imbalance²⁰⁴.

Controlled exposure to sodic-boric sodic layers and chilling conditions confirms that exogenous Si attenuates malondialdehyde (MDA) accumulation, an indicator of lipid peroxidation^{32,205}. Further, Liang et al.¹⁹⁶ reported that Si supplementation ameliorates freezing injury in overwinter wheat by remodeling the biophysical and biochemical attributes of the leaf apoplast, effectively substituting for the natural cold-acclimation response. Mechanistically, Si may fortify the antioxidative apparatus and stabilize the plasma membrane architecture, thereby diminishing cellular injury under diverse abiotic stress conditions¹³⁷. Support for agronomic applicability is evident; Datnoff et al.²⁰⁶ documented enhancement in growth metrics, biomass allocation, yield components and successful fertilization as a consequence of foliar and root Si feeding.

In many grape-growing locations, grape (*Vitis vinifera* L.) is vulnerable to cold temperatures²⁰⁷. Foliar Si treatment can successfully mitigate freezing damage in grapes by preserving membrane integrity and reducing photoinhibition during recovery²⁰⁸. Colder temperatures may lead to harm in pistachio growing areas, although foliar-applied Si mitigates the negative impacts of cold by raising Relative Water Content (RWC) and soluble sugar content in pistachio²⁰⁹. Cucumber is a vegetable crop that is vulnerable to chilling¹⁹². Exogenous Si treatment boosts endogenous

silicon deposition, boosts antioxidant activity and lowers lipid peroxidation caused by chilling stress in cucumber³². Cucumber cold resistance is improved by grafting and Si addition²¹⁰. Treatments such as salicylic acid or potassium silicate dipping keep cucumber fruits' freshness, flavour and appearance while also preventing decay and chilling damage symptoms²¹¹. Chitosan/nano-silica coating improves chilling resistance and increases loquat fruit storage life²¹². The addition of chitosan and nano-silica enhances the quality of longan fruit during storage at room temperature²¹³. Cold stress is a harmful abiotic stress that lowers medicinal plant output²¹⁴. Silicon has been shown to aid Sorghum bicolor growth under salt stress²¹⁵. Some wheat varieties thrive under cold conditions¹⁹⁶.

Silicon's role in improving oxidative stress tolerance: Exposure to abiotic stress markedly elevates the production of reactive oxygen species (ROS)-chiefly hydrogen peroxide (H_2O_2), superoxide ($O_2\cdot^-$) and hydroxyl radicals ($\cdot OH$)-within plant tissues, thereby threatening cellular integrity through oxidative damage to membranes and membranes and organelles²¹⁶. The plant antioxidant apparatus comprises both enzymatic and nonenzymatic components, the former including catalase (CAT), peroxidase (POD), superoxide dismutase (SOD) and ascorbate peroxidase (APX), while the latter is represented by vitamin E, ascorbic acid and Glutathione Reductase (GR)²¹⁷. Published studies underscore that the protective efficacy of these antioxidants is, to varying extents, modulated by silicon (Si) amendment. For instance, Si supplementation augmented the activities of CAT, SOD and GR in watermelon tissues, whereas APX activity remained constant⁸¹. Conversely, in cucumber-another model species-application of Si raised the activities of APX, SOD, glutathione peroxidase (GPX) and GR, while CAT activity showed no measurable variation²¹⁸. In tomato, exogenous Si not only curtailed the accumulation of H_2O_2 but also enhanced aquaporin-mediated water flux, thereby facilitating increased water uptake¹¹². These divergent biochemical modulations suggest the selective responsiveness of antioxidant mechanisms to Si, a noteworthy property that might be tailored to ameliorate oxidative and osmotic stress in species-specific contexts. Okra, grapes, wheat, rice and other plant species have shown similar outcomes^{219,220}. Depending on the plant species and Si content, the regulation scheme may vary. POD activity rose and MDA concentrations reduced with Si addition in *Glycyrrhiza uralensis*, whereas SOD activity increased only at a certain Si concentration²²¹.

Silicon influences plant antioxidant systems in a manner that is contingent upon species-specific responses, exposure duration, concentration and agronomic conditions, arguably succeeding in decreasing Reactive Oxygen Species (ROS) accumulation chiefly through the simultaneous regulation

of both enzymatic and non-enzymatic antioxidant mechanisms²²². Nevertheless, extensive agronomic trials remain imperative to ascertain the practical utility of silicon in commercial systems, since in vitro conditions generally fail to replicate the protracted and fluctuating abiotic stressors that characterize natural cropping environments. Interest in this regard has focused on the application of silicon compounds via foliar spray; evidence confirms that this method consistently enhances vegetative growth, elevates yield metrics and ameliorates stress symptoms across a broad taxonomic array of cultivated plant species²²³. Given that the silicon is made directly available to aboveground tissues, foliar application is postulated to confer a comparative advantage over traditional soil incorporation. Additionally, the direct supply of silicon to leaves may positively influence root morphology and length, consequently widening the effective root fringe and promoting improved nitrogen uptake, as observed in the recent comparative investigations by Kovács et al.²²⁴. Nonetheless, the extant literature does not provide a comprehensive, comparative evaluation of the relative efficiencies of soil- versus foliar-applied silicon; focused agronomic trials that systematically vary both application modes, concentrations and frequencies in parallel are prudent for fully characterizing the effective dosage and mode of delivery in differing cropping systems.

CONCLUSION

Silicon has been widely researched for its effect on plant developmental processes and physiological activities. It is necessary for maintaining ionic equilibrium, controlling water status, enabling photosynthesis, altering phytohormone levels and performing other important tasks. Furthermore, Si has been reported to promote antioxidative responses, raise osmolyte levels and stimulate stress-related signaling pathways in plants under stressful circumstances. Si is also effective in mitigating both biotic and abiotic stress. It increases the release of antimicrobial compounds within plants, assisting in pathogen defense and lowering infection rates. Furthermore, modern molecular approaches such as metabolomics, proteomics and transcriptomics have revealed on Si's exceptional significance in plants. Furthermore, research has shown that phytohormones crosstalk is involved, in which distinct hormones from plants and their signaling pathways interact in response to Si. However, more study is needed to comprehend the regulatory processes governing Si responses in plants, both in regular and stressful situations. Emphasizing genetic engineering methods can aid in the discovery of metabolic pathways responsible for various plant characteristics connected to Si. We can acquire a more thorough grasp of Si's impacts on plant physiology and its potential for stress management by going deeper into these areas.

REFERENCES

1. Kostina-Bednarz M, Plonka J, Barchanska H. Allelopathy as a source of bioherbicides: Challenges and prospects for sustainable agriculture. *Rev Environ Sci Bio/Technol.* 2023;22:471-504.
2. Etesami H, Jeong BR. Silicon (Si): Review and future prospects on the action mechanisms in alleviating biotic and abiotic stresses in plants. *Ecotoxicol Environ Saf.* 2018;147:881-896.
3. Malhi GS, Kaur M, Kaushik P. Impact of climate change on agriculture and its mitigation strategies: A review. *Sustainability.* [Internet]. 2021;13(3). Available from: <https://doi.org/10.3390/su13031318>
4. Sahebi M, Hanafi MM, Akmar ASN, Rafii MY, Azizi P, Tengoua FF et al. Importance of silicon and mechanisms of biosilica formation in plants. *Biomed Res Int.* [Internet]. 2015;2015. Available from: <https://doi.org/10.1155/2015/396010>
5. Minhas PS, Rane J, Pasala RK. Abiotic stress management for resilient agriculture. 1st ed. Singapore: Springer Singapore; 2017.
6. Santos TBD, Ribas AF, Souza SGHD, Budzinski IGF, Domingues DS. Physiological responses to drought, salinity and heat stress in plants: A review. *Stresses.* [Internet]. 2022;2(1). Available from: <https://doi.org/10.3390/stresses2010009>
7. Berg G. Plant-microbe interactions promoting plant growth and health: Perspectives for controlled use of microorganisms in agriculture. *Appl Microbiol Biotechnol.* 2009;84:11-18.
8. Schmidt R, Köberl M, Mostafa A, Ramadan EM, Monschein M, Jensen KB et al. Effects of bacterial inoculants on the indigenous microbiome and secondary metabolites of chamomile plants. *Front Microbiol.* [Internet]. 2014;5. Available from: <https://doi.org/10.3389/fmicb.2014.00064>
9. Oyedoh OP, Yang W, Dhanasekaran D, Santoyo G, Glick BR, Babalola OO. Rare rhizo-actinomycetes: A new source of agroactive metabolites. *Biotechnol Adv.* [Internet]. 2023;67. Available from: <https://doi.org/10.1016/j.biotechadv.2023.108205>
10. Kumari P, Meena M, Gupta P, Dubey MK, Nath G, Upadhyay R. Plant growth promoting rhizobacteria and their bioprimeing for growth promotion in mung bean (*Vigna radiata* (L.) R. Wilczek). *Biocatal Agric Biotechnol.* 2018;16:163-171.
11. Son JS, Sumayo M, Hwang YJ, Kim BS, Ghim SY. Screening of plant growth-promoting rhizobacteria as elicitor of systemic resistance against gray leaf spot disease in pepper. *Appl Soil Ecol.* 2014;73:1-8.
12. Zargar SM, Mahajan R, Bhat JA, Nazir M, Deshmukh R. Role of silicon in plant stress tolerance: Opportunities to achieve a sustainable cropping system. *3 Biotech.* [Internet]. 2019;9. Available from: <https://doi.org/10.1007/s13205-019-1613-z>
13. Ma JF. Role of silicon in enhancing the resistance of plants to biotic and abiotic stresses. *Soil Sci Plant Nutr.* 2004;50(1):11-18.
14. Stein M, Buchweitz M, Mayer P, Rennert T. Does adsorption of Cd, Cu and Pb on polymeric silicic acid occur under acidic conditions? *Silicon.* 2023;15:7205-7212.
15. Savvas D, Ntatsi G. Biostimulant activity of silicon in horticulture. *Sci Hortic.* 2015;196:66-81.
16. Farooq MA, Dietz KJ. Silicon as versatile player in plant and human biology: Overlooked and poorly understood. *Front Plant Sci.* [Internet]. 2015;6. Available from: <https://doi.org/10.3389/fpls.2015.00994>
17. Iezzana T, Sigler K. Biologically active compounds of semi-metals. *Stud Nat Prod Chem.* 2008;35:835-921.
18. Raza A. Eco-physiological and biochemical responses of rapeseed (*Brassica napus* L.) to abiotic stresses: Consequences and mitigation strategies. *J Plant Growth Regul.* 2020;40:1368-1388.
19. Thakral V, Bhat JA, Kumar N, Myaka B, Sudhakaran S, Patil G et al. Role of silicon under contrasting biotic and abiotic stress conditions provides benefits for climate smart cropping. *Environ Exp Bot.* [Internet]. 2021;189. Available from: <https://doi.org/10.1016/j.envexpbot.2021.104545>
20. Munir FDA, Sulaiman Z, Ahmad K, Samad MYA, Ariffin MR, Salisu MA. A comprehensive review on the potential effect of silicon fertilizer and its significance for agriculture. *Icrrd Qual Index Res J.* [Internet]. 2023;4. Available from: <https://doi.org/10.53272/icrrd.v4i2.5>
21. Ali M, Afzal S, Parveen A, Kamran M, Javed MR, Abbasi GH et al. Silicon mediated improvement in the growth and ion homeostasis by decreasing Na^+ uptake in maize (*Zea mays* L.) cultivars exposed to salinity stress. *Plant Physiol Biochem.* 2021;158:208-218.
22. Shen Z, Pu X, Wang S, Dong X, Cheng X, Cheng M. Silicon improves ion homeostasis and growth of liquorice under salt stress by reducing plant Na^+ uptake. *Sci Rep.* [Internet]. 2022;12. Available from: <https://doi.org/10.1038/s41598-022-09061-8>
23. Tripathi DK, Vishwakarma K, Singh VP, Prakash V, Sharma S, Muneer S et al. Silicon crosstalk with reactive oxygen species, phytohormones and other signaling molecules. *JHazard Mater.* [Internet]. 2021;408. Available from: <https://doi.org/10.1016/j.jhazmat.2020.124820>
24. Bhardwaj S, Sharma D, Singh S, Ramamurthy PC, Verma T, Pujari M et al. Physiological and molecular insights into the role of silicon in improving plant performance under abiotic stresses. *Plant Soil.* 2022;486:25-43.
25. Abbas T, Balal RM, Shahid MA, Pervez MA, Ayyub CM, Aqueel MA et al. Silicon-induced alleviation of NaCl toxicity in okra (*Abelmoschus esculentus*) is associated with enhanced photosynthesis, osmoprotectants and antioxidant metabolism. *Acta Physiol Plant.* [Internet]. 2015;37. Available from: <https://doi.org/10.1007/s11738-014-1768-5>
26. Zhu Y, Gong H. Beneficial effects of silicon on salt and drought tolerance in plants. *Agron Sustainable Dev.* 2013;34:455-472.

27. Zhu YX, Gong HJ, Yin JL. Role of silicon in mediating salt tolerance in plants: A review. *Plants*. [Internet]. 2019;8(6). Available from: <https://doi.org/10.3390/plants8060147>
28. Artyszak A. Effect of silicon fertilization on crop yield quantity and quality-A literature review in Europe. *Plants*. [Internet]. 2018;7(3). Available from: <https://doi.org/10.3390/plants7030054>
29. Soppelsa S, Kelderer M, Casera C, Bassi M, Robatscher P and reotti C. Use of biostimulants for organic apple production: Effects on tree growth, yield and fruit quality at harvest and during storage. *Front Plant Sci*. [Internet]. 2018;9. Available from: <https://doi.org/10.3389/fpls.2018.01342>
30. Ahmad S, Pathak AK, Minj RP, Chaudhary S, Kumar A, Chalotra T et al. A comprehensive review of nanoparticles induced stress and toxicity in plants. *Universal J Green Chem*. 2023;1(1):18-43.
31. Mir RA, Bhat BA, Yousuf H, Islam ST, Raza A, Rizvi MA et al. Multidimensional role of silicon to activate resilient plant growth and to mitigate abiotic stress. *Front Plant Sci*. [Internet]. 2022;13. Available from: <https://doi.org/10.3389/fpls.2022.819658>
32. LIU JJ, Lin SH, XU PL, Wang XJ, Bai JG. Effects of exogenous silicon on the activities of antioxidant enzymes and lipid peroxidation in chilling-stressed cucumber leaves. *Agric Sci China*. 2009;8(9):1075-1086.
33. Silva MND, Machado J, Osorio J, Duarte R, Santos CS. Non-essential elements and their role in sustainable agriculture. *Agronomy*. [Internet]. 2022;12(4). Available from: <https://doi.org/10.3390/agronomy12040888>
34. Fellet G, Pilotto L, Marchiol L, Braidot E. Tools for nano-enabled agriculture: Fertilizers based on calcium phosphate, silicon and chitosan nanostructures. *Agronomy*. [Internet]. 2021;11(6). Available from: <https://doi.org/10.3390/agronomy11061239>
35. Liang Y, Nikolic M, Bélanger R, Gong H; Song A. Silicon-mediated tolerance to metal toxicity. In: Liang Y, Nikolic M, Bélanger R, Gong H, Song A, eds. *Silicon in Agriculture*. Springer Netherlands; 2015:83-122. 10.1007/978-94-017-9978-2_5
36. Singh A, Roychoudhury A. Abscisic acid in plants under abiotic stress: Crosstalk with major phytohormones. *Plant Cell Rep*. 2023;42:961-974.
37. Cheng Z, Song W, Zhang X. Genic male and female sterility in vegetable crops. *Hortic Res*. [Internet]. 2022;10(1). Available from: <https://doi.org/10.1093/hr/uhac232>
38. Al-Khayri JM, Rashmi R, Ulhas RS, Sudheer WN, Banadka A, Nagella P et al. The role of nanoparticles in response of plants to abiotic stress at physiological, biochemical and molecular levels. *Plants*. [Internet]. 2023;12(2). Available from: <https://doi.org/10.3390/plants12020292>
39. Mostofa MG, Rahman M, Ansary MU, Keya SS, Abdelrahman M, Miah G et al. Silicon in mitigation of abiotic stress-induced oxidative damage in plants. *Crit Rev Biotechnol*. 2021;41(6):918-934.
40. Hasanuzzaman M, Bhuyan M, Zulfiqar F, Raza A, Mohsin S, Mahmud J et al. Reactive oxygen species and antioxidant defense in plants under abiotic stress: Revisiting the crucial role of a universal defense regulator. *Antioxidants*. [Internet]. 2020;9(8). Available from: <https://doi.org/10.3390/antiox9080681>
41. Swain R, Sahoo S, Behera M, Rout GR. Instigating prevalent abiotic stress resilience in crop by exogenous application of phytohormones and nutrient. *Front Plant Sci*. [Internet]. 2023;14. Available from: <https://doi.org/10.3389/fpls.2023.1104874>
42. Avasiloaiei DI, Calara M, Brezeanu PM, Murariu OC, Brezeanu C. On the future perspectives of some medicinal plants within Lamiaceae botanic family regarding their comprehensive properties and resistance against biotic and abiotic stresses. *Genes*. [Internet]. 2023;14(5). Available from: <https://doi.org/10.3390/genes14050955>
43. El-Okkiah SAF, El-Tahan AM, Ibrahim OM, Taha MA, Korany SM, Alsherif EA et al. Under cadmium stress, silicon has a defensive effect on the morphology, physiology and anatomy of pea (*Pisum sativum L.*) plants. *Front Plant Sci*. [Internet]. 2022;13. Available from: <https://doi.org/10.3389/fpls.2022.997475>
44. Siddiqui H, Yusuf M, Faraz A, Faizan M, Sami F, Hayat S. 24-epibrassinolide supplemented with silicon enhances the photosynthetic efficiency of *Brassica juncea* under salt stress. *South Afr J Bot*. 2018;118(118):120-128.
45. Patel M, Fatnani D, Parida AK. Silicon-induced mitigation of drought stress in peanut genotypes (*Arachis hypogaea L.*) through ion homeostasis, modulations of antioxidative defense system and metabolic regulations. *Plant Physiol Biochem*. 2021;166:290-313.
46. Kim YH, Khan AL, Waqas M, Jeong HJ, Kim DH, Shin JS et al. Regulation of jasmonic acid biosynthesis by silicon application during physical injury to *Oryza sativa L.* *J Plant Res*. 2014;127:525-532.
47. Sabagh AE, Islam MS, Hossain A, Iqbal MA, Mubeen M, Waleed M et al. Phytohormones as growth regulators during abiotic stress tolerance in plants. *Front Agr*. [Internet]. 2022;4. Available from: <https://doi.org/10.3389/fagro.2022.765068>
48. Sharma A, Sidhu GPS, Araniti F, Bali AS, Shahzad B, Tripathi DK et al. The role of salicylic acid in plants exposed to heavy metals. *Molecules*. [Internet]. 2020;25(3). Available from: <https://doi.org/10.3390/molecules25030540>
49. Bharath P, Gahir S, Raghavendra AS. Abscisic acid-induced stomatal closure: An important component of plant defense against abiotic and biotic stress. *Front Plant Sci*. [Internet]. 2021;12. Available from: <https://doi.org/10.3389/fpls.2021.615114>

50. Wahab A, Abdi G, Saleem MH, Ali B, Ullah S, Shah W et al. Plants' physio-biochemical and phytohormonal responses to alleviate the adverse effects of drought stress: A comprehensive review. *Plants*. [Internet]. 2022;11(13). Available from: <https://doi.org/10.3390/plants11131620>

51. Xu BQ, Wang JJ, Peng Y, Huang H, Sun LL, Yang R et al. SIMYC2 mediates stomatal movement in response to drought stress by repressing S1CHS1 expression. *Front Plant Sci*. [Internet]. 2022;13. Available from: <https://doi.org/10.3389/fpls.2022.952758>

52. Kumar S, Shah SH, Vimala Y, Jatav HS, Ahmad P, Chen Y et al. Abscisic acid: Metabolism, transport, crosstalk with other plant growth regulators and its role in heavy metal stress mitigation. *Front Plant Sci*. [Internet]. 2022;13. Available from: <https://doi.org/10.3389/fpls.2022.972856>

53. Liu B, Soundararajan P, Manivannan A. Mechanisms of silicon-mediated amelioration of salt stress in plants. *Plants*. [Internet]. 2019;8(9). Available from: <https://doi.org/10.3390/plants8090307>

54. Khan MIR, Ashfaque F, Chhillar H, Irfan M, Khan NA. The intricacy of silicon, plant growth regulators and other signaling molecules for abiotic stress tolerance: An entrancing crosstalk between stress alleviators. *Plant Physiol Biochem*. 2021;162:36-47.

55. Nguyen HT, Leipner J, Stamp P, Guerra-Peraza O. Low temperature stress in maize (*Zea mays* L.) induces genes involved in photosynthesis and signal transduction as studied by suppression subtractive hybridization. *Plant Physiol Biochem*. 2009;47(2):116-122.

56. Gill SS, Tuteja N. Polyamines and abiotic stress tolerance in plants. *Plant Signaling Behav*. 2010;5(1):26-33.

57. Ahanger MA, Bhat JA, Siddiqui MH, Rinklebe J, Ahmad P. Integration of silicon and secondary metabolites in plants: A significant association in stress tolerance. *J Exp Bot*. 2020;71(21):6758-6774.

58. El-Serafy RS, El-Sheshtawy ANA, Atteya AK, Al-Hashimi A, Abbasi AM, Al-Ashkar I. Seed priming with silicon as a potential to increase salt stress tolerance in *Lathyrus odoratus*. *Plants*. [Internet]. 2021;10(10). Available from: <https://doi.org/10.3390/plants10102140>

59. Wang Z, Poudyal S, Kopp K, Zhang Y. Response of ornamental plants to salinity: Impact on species-specific growth, visual quality, photosynthetic parameters and ion uptake. *Front Plant Sci*. [Internet]. 2025;16. Available from: <https://doi.org/10.3389/fpls.2025.1611767>

60. Hasan K, Sabagh AE, Sikdar SI, Alam J, Ratnasekera D, Barutcular C et al. Comparative adaptable agronomic traits of blackgram and mungbean for saline lands. *Plant Arch*. 2017;17(1):589-593.

61. Agarie S, Hanaoka N, Ueno O, Miyazaki A, Kubota F, Agata W et al. Effects of silicon on tolerance to water deficit and heat stress in rice plants (*Oryza sativa* L.), monitored by electrolyte leakage. *Plant Prod Sci*. 1998;1(2):96-103.

62. Avestan S, Ghasemnezhad M, Esfahani M, Byrt CS. Application of nano-silicon dioxide improves salt stress tolerance in strawberry plants. *Agronomy*. [Internet]. 2019;9(5). Available from: <https://doi.org/10.3390/agronomy9050246>

63. Flam-Shepherd R, Huynh WQ, Coskun D, Hamam AM, Britto DT, Kronzucker HJ. Membrane fluxes, bypass flows and sodium stress in rice: The influence of silicon. *J Exp Bot*. 2018;69(7):1679-1692.

64. Bao-shan L, Shao-qi D, Chun-hui L, Li-jun F, Shu-chun Q, Min Y. Effect of TMS (nanostructured silicon dioxide) on growth of Changbai larch seedlings. *J For Res*. 2004;15:138-140.

65. Alva AK, Mattos D, Paramasivam S, Patil B, Dou H, Sajwan KS. Potassium management for optimizing citrus production and quality. *Int J Fruit Sci*. 2006;6(1):3-43.

66. Almeida DM, Oliveira MM, Saibo NJM. Regulation of Na^+ and K^+ homeostasis in plants: Towards improved salt stress tolerance in crop plants. *Genet Mol Biol*. 2017;40:326-345.

67. Rizwan M, Ali S, Ibrahim M, Farid M, Adrees M, Bharwana SA et al. Mechanisms of silicon-mediated alleviation of drought and salt stress in plants: A review. *Envir Sci Polluti Res*. 2015;22:15416-15431.

68. Seal P, Das P, Biswas AK. Versatile potentiality of silicon in mitigation of biotic and abiotic stresses in plants: A review. *Am J Plant Sci*. 2018;9(7):1433-1454.

69. Ahanger MA, Alyemeni MN, Wijaya L, Alamri SA, Alam P, Ashraf M et al. Potential of exogenously sourced kinetin in protecting *Solanum lycopersicum* from NaCl -induced oxidative stress through up-regulation of the antioxidant system, ascorbate-glutathione cycle and glyoxalase system. *PLoS ONE*. [Internet]. 2018;13. Available from: <https://doi.org/10.1371/journal.pone.0202175>

70. Ahmad P, Ahanger MA, Alam P, Alyemeni MN, Wijaya L, Ali S et al. Silicon (Si) supplementation alleviates NaCl toxicity in mung bean [*Vigna radiata* (L.) Wilczek] through the modifications of physio-biochemical attributes and key antioxidant enzymes. *J Plant Growth Regul*. 2018;38:70-82.

71. Sabaghnia N, Janmohammadi M. Effect of nano-silicon particles application on salinity tolerance in early growth of some lentil genotypes. *Ann Universitatis Mariae Curie-Sklodowska, sectio C-Biol*. 2015;62(2):39-55.

72. Sanzari I, Leone A, Ambrosone A. Nanotechnology in plant science: To make a long story short. *Front Bioeng Biotechnol*. [Internet]. 2019;7. Available from: <https://doi.org/10.3389/fbioe.2019.00120>

73. Zahedi SM, Karimi M, Silva JATD. The use of nanotechnology to increase quality and yield of fruit crops. *J Sci Food Agric*. 2019;100:25-31.

74. Abdelaal KAA, Mazrou YS, Hafez YM. Silicon foliar application mitigates salt stress in sweet pepper plants by enhancing water status, photosynthesis, antioxidant enzyme activity and fruit yield. *Plants*. [Internet]. 2020;9(6). Available from: <https://doi.org/10.3390/plants9060733>

75. Shi Y, Zhang Y, Yao H, Wu J, Sun H, Gong H. Silicon improves seed germination and alleviates oxidative stress of bud seedlings in tomato under water deficit stress. *Plant Physiol Biochem*. 2014;78:27-36.

76. Karunakaran G, Suriyaprabha R, Manivasakan P, Yuvakkumar R, Rajendran V, Prabu P et al. Effect of nanosilica and silicon sources on plant growth promoting rhizobacteria, soil nutrients and maize seed germination. *IET Nanobiotechnol*. 2013;7:70-77.

77. Islam MJ, Ryu BR, Azad MOK, Rahman MH, Rana MS, Lim JD et al. Exogenous putrescine enhances salt tolerance and ginsenosides content in Korean ginseng (*Panax ginseng* Meyer) sprouts. *Plants*. [Internet]. 2021;10(7). Available from: <https://doi.org/10.3390/plants10071313>

78. Pinedo-Guerrero ZH, Cadenas-Pliego G, Ortega-Ortiz H, González-Morales S, Benavides-Mendoza A, Valdés-Reyna J et al. Form of silica improves yield, fruit quality and antioxidant defense system of tomato plants under salt stress. *Agric*. [Internet]. 2020;10(9). Available from: <https://doi.org/10.3390/agriculture10090367>

79. Tantawy A, Salama Y, El-Nemr M, Abdel-Mawgoud A. Nano silicon application improves salinity tolerance of sweet pepper plants. *Int J Chemtech Res*. 2015;8(10):11-17.

80. Elsheery NI, Helaly MN, El-Hoseiny HM, Alam-Eldein SM. Zinc oxide and silicone nanoparticles to improve the resistance mechanism and annual productivity of salt-stressed mango trees. *Agronomy*. [Internet]. 2020;10(4). Available from: <https://doi.org/10.3390/agronomy10040558>

81. Bijalwan P, Jreddi K, Saini I, Sharma M, Kaushik P, Hessini K. Mitigation of saline conditions in watermelon with mycorrhiza and silicon application. *Saudi J Biol Sci*. 2021;28(7):3678-3684.

82. Patil H, Tank RV, Manoli P. Significance of silicon in fruit crops - A review. *Plant Arch*. 2017;17(2):769-774.

83. Li H, Zhu Y, Hu Y, Han W, Gong H. Beneficial effects of silicon in alleviating salinity stress of tomato seedlings grown under sand culture. *Acta Physiol Plant*. [Internet]. 2015;37. Available from: <https://doi.org/10.1007/s11738-015-1818-7>

84. Muneer S, Jeong BR. Proteomic analysis of salt-stress responsive proteins in roots of tomato (*Lycopersicon esculentum* L.) plants towards silicon efficiency. *Plant Growth Regul*. 2015;77:133-146.

85. Lee SK, Sohn EY, Hamayun M, Yoon JY, Lee IJ. Effect of silicon on growth and salinity stress of soybean plant grown under hydroponic system. *Agrofor Syst*. 2010;80:333-340.

86. Samuels AL, Glass AD, Ehret DL, Menzies JG. The effects of silicon supplementation on cucumber fruit: Changes in surface characteristics. *Ann Bot*. 1993;72(5):433-440.

87. Ullah H, Santiago-Arenas R, Ferdous Z, Attia A, Datta A. Improving water use efficiency, nitrogen use efficiency and radiation use efficiency in field crops under drought stress: A review. In: Sparks DL, ed. *Advances in Agronomy*. United States: Academic Press; 2019. 109-157.

88. Sirisuntornlak N, Ghafoori S, Datta A, Arirob W. Seed priming and soil incorporation with silicon influence growth and yield of maize under water-deficit stress. *Arch Agron Soil Sci*. 2018;65:197-207.

89. Farooq M, Wahid A, Kobayashi N, Fujita D, Basra SMA. Plant drought stress: Effects, mechanisms and management. *Agron Sustainable Dev*. 2009;29:185-212.

90. Wang M, Wang R, Mur LAJ, Ruan J, Shen Q, Guo S. Functions of silicon in plant drought stress responses. *Hortic Res*. [Internet]. 2021;8. Available from: <https://doi.org/10.1038/s41438-021-00681-1>

91. Bukhari MA, Ashraf MY, Ahmad R, Waraich EA, Hameed M. Improving drought tolerance potential in wheat (*Triticum aestivum* L.) Through exogenous silicon supply. *Pak J Bot*. 2015;47(5):1641-1648.

92. Sayfzadeh S, Rashidi M. Response of antioxidant enzyme activities of sugar beet to drought stress. *ARPV J Agric Biol Sci*. 2011;16(4):27-33.

93. Cifre J, Bota J, Escalona J, Medrano H, Flexas J. Physiological tools for irrigation scheduling in grapevine (*Vitis vinifera* L.): An open gate to improve water-use efficiency? *Agric, EcoSyst Environ*. 2005;106:159-170.

94. Masoumi Z, Haghghi M, Mozafarian M. Effects of foliar spraying with melatonin and chitosan Nano-encapsulated melatonin on tomato (*Lycopersicon esculentum* L. cv. Falcato) plants under salinity stress. *BMC Plant Biol*. [Internet]. 2024;24. Available from: <https://doi.org/10.1186/s12870-024-05672-7>

95. Schieber M, Chandel N. Ros function in redox signaling and oxidative stress. *Curr Biol*. 2014;24(10):453-462.

96. Cotrim CE, Filho MAC, Coelho EF, Ramos MM, Cecon PR. Regulated deficit irrigation and Tommy Atkins mango orchard productivity under microsprinkling in Brazilian semi arid. *Eng Agric*. 2011;31(6):1052-1063.

97. Mishra N, Jiang C, Chen L, Paul A, Chatterjee A, Shen G. Achieving abiotic stress tolerance in plants through antioxidative defense mechanisms. *Front Plant Sci*. [Internet]. 2023;14. Available from: <https://doi.org/10.3389/fpls.2023.1110622>

98. Balakhnina T, Borkowska A. Effects of silicon on plant resistance to environmental stresses: Review. *Int Agrophys*. 2013;27:225-232.

99. Cattivelli L, Rizza F, Badeck FW, Mazzucotelli E, Mastrangeli AM, Francia E et al. Drought tolerance improvement in crop plants: An integrated view from breeding to genomics. *Field Crops Res*. 2008;105:1-14.

100. Sharma SP, Leskovar DI, Crosby KM, Volder A, Ibrahim A. Root growth, yield and fruit quality responses of reticulatus and inodorus melons (*Cucumis melo* L.) to deficit subsurface drip irrigation. *Agric Water Manage.* 2014;136:75-85.

101. Mauad M, Cruscio CAC, Nascente AS, Filho HG, Lima GPP. Effects of silicon and drought stress on biochemical characteristics of leaves of upland rice cultivars. *Rev Cienc Agron.* 2016;47(3):532-539.

102. Neeru J, Shaliesh C, Vaishali T, Purav S, Manoharlal R. Role of orthosilicic acid (OSA) based formulation in improving plant growth and development. *Silicon.* 2016;11:2407-2411.

103. El-Shazoly RM. Ameliorating effect of silicon on growth vigor, physiological and biochemical traits of salinized canola seedlings (*Brassica napus* L.). *Catrina: Int J Environ Sci.* 2019;18(1):125-137.

104. Nascimento CWAD, Nunes GHDS, Preston HAF, Silva FBVD, Preston W, Loureiro FLC. Influence of silicon fertilization on nutrient accumulation, yield and fruit quality of melon grown in northeastern Brazil. *Silicon.* 2019;12:937-943.

105. Toresano-Sánchez F, Valverde-García A, Camacho-Ferre F. Effect of the application of silicon hydroxide on yield and quality of cherry tomato. *J Plant Nutr.* 2012;35(4):567-590.

106. Kleiber T, Kleiber T, Bosiacki M, Brell W. Effect of choline-stabilized orthosilic acid application on tomato grown under increasing Mn stress. *J Elementology.* 2015;20(4):897-910.

107. Greger M, Landberg T, Vaculík M. Silicon influences soil availability and accumulation of mineral nutrients in various plant species. *Plants.* [Internet]. 2018;7(2). Available from: <https://doi.org/10.3390/plants7020041>

108. Alam A, Hariyanto B, Ullah H, Salin KR, Datta A. Effects of silicon on growth, yield and fruit quality of cantaloupe under drought stress. *Silicon.* 2020;13:3153-3162.

109. Zahedi SM, Moharrami F, Sarikhani S, Padervand M. Selenium and silica nanostructure-based recovery of strawberry plants subjected to drought stress. *Sci Rep.* [Internet]. 2020;10. Available from: <https://doi.org/10.1038/s41598-020-74273-9>

110. Helaly MN, El-Hoseiny H, El-Sheery NI, Rastogi A, Kalaji HM. Regulation and physiological role of silicon in alleviating drought stress of mango. *Plant Physiol Biochem.* 2017;118:31-44.

111. Tabrizian ST, Hajilou J, Bolandnazar S, Dehghan G. Silicon improves strawberry ability to cope with water deficit stress. *Int J Hortic Sci Technol.* 2020;9(2):213-226.

112. Shi Y, Zhang Y, Han W, Feng R, Hu Y, Guo J et al. Silicon enhances water stress tolerance by improving root hydraulic conductance in *Solanum lycopersicum* L.. *Front Plant Sci.* [Internet]. 2016;7. Available from: <https://doi.org/10.3389/fpls.2016.00196>

113. Cantore V, Lechkar O, Karabulut E, Sellami MH, Albrizio R, Boari F et al. Combined effect of deficit irrigation and strobilurin application on yield, fruit quality and water use efficiency of "cherry" tomato (*Solanum lycopersicum* L.). *Agric Water Manage.* 2016;167:53-61.

114. Chakma R, Saekong P, Biswas A, Ullah H, Datta A. Growth, fruit yield, quality and water productivity of grape tomato as affected by seed priming and soil application of silicon under drought stress. *Agric Water Manage.* [Internet]. 2021;256. Available from: <https://doi.org/10.1016/j.agwat.2021.107055>

115. Aziz MAE. The effect of si on minimizing the implications of water stress on tomato plants. *Environ, Biodivers Soil Secur.* 2020;4:137-148.

116. Maestro MC, Alvarez J. The effects of temperature on pollination and pollen tube growth in muskmelon (*Cucumis melo* L.). *Sci Hortic.* 1988;36:173-181.

117. El-Kader AMA, Saleh MMS, Ali MA. Effect of soil moisture levels and some antitranspirants on vegetative growth, leaf mineral content, yield and fruit quality of Williams banana plants. *J Appl Sci Res.* 2006;2(12):1248-1255.

118. Thornton I. Metal contamination of soils in urban areas. In: Bullock P, Gregory PJ, eds. *Soils in the Urban Environment.* Oxford: Wiley; 1991. 47-75.

119. Chen Z, Muhammad I, Zhang Y, Hu W, Lu Q, Wang W et al. Transfer of heavy metals in fruits and vegetables grown in greenhouse cultivation systems and their health risks in Northwest China. *Sci Total Environ.* [Internet]. 2021;766. Available from: <https://doi.org/10.1016/j.scitotenv.2020.142663>

120. Meena V, Dotaniya ML, Saha JK, Patra AK. Silicon potential to mitigate plant heavy metals stress for sustainable agriculture: A review. *Silicon.* 2021;14:4447-4462.

121. Stewart H, Hyman J, Dong D, Carlson A. The more that households prioritise healthy eating, the better they can afford to consume a sufficient quantity and variety of fruits and vegetables. *Public Health Nutr.* 2020;24(7):1841-1850.

122. Zwolak A, Sarzyńska M, Szpyrka E, Stawarczyk K. Sources of soil pollution by heavy metals and their accumulation in vegetables: A review. *Water, Air, Soil Pollut.* [Internet]. 2019;230. Available from: <https://doi.org/10.1007/s11270-019-4221-y>

123. Wu L, Sun Q, Zhao JM, Wang DL, Zhang YW. Review on the effects of heavy metal accumulation in flowers on the mutual benefit relationship between plant and insect pollinators. *Ying Yong Sheng Tai Xue Bao = J Appl Ecol.* 2022;33(5):1429-1429.

124. Guo C, Lv L, Liu Y, Ji M, Zang E, Liu Q et al. Applied analytical methods for detecting heavy metals in medicinal plants. *Crit Revs Anal Chem.* 2021;53(2):339-359.

125. Liang G, Gong W, Li B, Zuo J, Pan L, Liu X. Analysis of heavy metals in foodstuffs and an assessment of the health risks to the general public via consumption in Beijing, China. *Int J Environ Res Public Health*. [Internet]. 2019;16(6). Available from: <https://doi.org/10.3390/ijerph16060909>

126. Kim Y, Ha EH, Park H, Ha M, Kim Y, Hong YC et al. Prenatal lead and cadmium co-exposure and infant neurodevelopment at 6 months of age: The mothers and children's environmental health (MOCEH) study. *NeuroToxicology*. 2013;35:15-22.

127. Reuben A, Caspi A, Belsky DW, Broadbent J, Harrington H, Sugden K et al. Association of childhood blood lead levels with cognitive function and socioeconomic status at age 38 years and with IQ change and socioeconomic mobility between childhood and adulthood. *JAMA*. 2017;317(12):1244-0.

128. Mawari G, Kumar N, Sarkar S, Daga MK, Singh MM, Joshi TK et al. Heavy metal accumulation in fruits and vegetables and human health risk assessment: Findings from Maharashtra, India. *Environ Health Insights*. [Internet]. 2022;16. Available from: <https://doi.org/10.1177/11786302221119151>

129. Gudadhe S, Singh SK, Ahsan J. Cellular and neurological effects of lead (Pb) toxicity. In: Kumar N, Jha AK, eds. *Lead Toxicity Mitigation: Sustainable Nexus Approaches*. Cham: Springer Nature Switzerland; 2024. 125-145.

130. Dotaniya ML, Dotaniya CK, Solanki P, Meena VD, Doutaniya RK. Lead contamination and its dynamics in soil-Plant system. In: Gupta DK, Chatterjee S, Walther C, eds. *Lead in Plants and the Environment*. Cham: Springer International Publishing; 2019. 83-98.

131. Vasilachi IC, Stoleru V, Gavrilescu M. Analysis of heavy metal impacts on cereal crop growth and development in contaminated soils. *Agricultur*. [Internet]. 2023;13(10). Available from: <https://doi.org/10.3390/agriculture13101983>

132. Islam MS, Ahmed MK, Habibullah-Al-Mamun M, Raknuzzaman M, Ali MM, Eaton DW. Health risk assessment due to heavy metal exposure from commonly consumed fish and vegetables. *Environ Syst Decisions*. 2016;36:253-265.

133. Li Q, Cai S, Mo C, Chu B, Peng L, Yang F. Toxic effects of heavy metals and their accumulation in vegetables grown in a saline soil. *Ecotoxicol Environ Saf*. 2010;73(1):84-88.

134. Kumari VV, Roy A, Vijayan R, Banerjee P, Verma VC, Nalia A et al. Drought and heat stress in cool-season food legumes in sub-tropical regions: Consequences, adaptation and mitigation strategies. *Plants*. [Internet]. 2021;10(6). Available from: <https://doi.org/10.3390/plants10061038>

135. LİCİTUŞU R, LİCİTUŞU AR. Vegetable and fruits quality within heavy metals polluted areas in Romania. *Carpathian J Earth Environ Sci*. 2008;3(2):115-129.

136. Gong Hj, Chen Km, Wang Sm. Advances in silicon nutrition of plants. *Acta Botanica Boreali-Occidentalia Sin*. [Internet]. 2004;24(12). Available from: <http://xbzwxb.alljournal.net/xbzwxben/article/abstract/2004012520>

137. Liang Y, Sun W, Zhu YG, Christie P. Mechanisms of silicon-mediated alleviation of abiotic stresses in higher plants: A review. *Environ Pollut*. 2007;147(2):422-428.

138. Rodrigues FA, Datnoff LE. *Silicon and plant diseases*. 1st ed. Cham, Switzerland: Springer International Publishing; 2015.

139. Emamverdian A, Ding Y, Xie Y, Sangari S. Silicon mechanisms to ameliorate heavy metal stress in plants. *Biomed Res Int*. [Internet]. 2018. Available from: <https://doi.org/10.1155/2018/8492898>

140. Kim YH, Khan AL, Kim DH, Lee SY, Kim KM, Waqas M et al. Silicon mitigates heavy metal stress by regulating P-type heavy metal ATPases, *Oryza satival*ow silicon genes and endogenous phytohormones. *BMC Plant Biol*. [Internet]. 2014;14. Available from: <https://doi.org/10.1186/1471-2229-14-13>

141. Wu J, Mock HP, Giehl RF, Pitann B, Mühling KH. Silicon decreases cadmium concentrations by modulating root endodermal suberin development in wheat plants. *J Hazard Mater*. 2019;364:581-590.

142. Rachappanavar V, Gupta SK, Jayaprakash GK, Abbas M. Silicon mediated heavy metal stress amelioration in fruit crops. *Helijon*. [Internet]. 2024;10(18). Available from: <https://doi.org/10.1016/j.helijon.2024.e37425>

143. Stadnik B, Tobiasz-Salach R, Mazurek M. Physiological and epigenetic reaction of barley (*Hordeum vulgare L.*) to the foliar application of silicon under soil salinity conditions. *Int J Mol Sci*. [Internet]. 2022;23(3). Available from: <https://doi.org/10.3390/ijms23031149>

144. Debona D, Rodrigues FA, Datnoff LE. Silicon's role in abiotic and biotic plant stresses. *Annu Rev Phytopathol*. 2017;55:85-107.

145. Hodson MJ, White PJ, Mead A, Broadley MR. Phylogenetic variation in the silicon composition of plants. *Ann Bot*. 2005;96(6):1027-1046.

146. Ma JF, Yamaji N. Silicon uptake and accumulation in higher plants. *Trends Plant Sci*. 2006;11(8):392-397.

147. Wedepohl KH. The composition of the continental crust. *Geochim Cosmochim Acta*. 1995;59(7):1217-1232.

148. Kowalska J, Tyburski J, Jakubowska M, Krzymińska J. Effect of different forms of silicon on growth of spring wheat cultivated in organic farming system. *Silicon*. 2020;13:211-217.

149. Szymalska R. *Responses of Plants to Environmental Stresses*. Basel, Switzerland: MDPI; 2021.

150. Yang C, Xu Y, Xie X, Wu Y, Gao Z, Li K et al. Post-harvest physiology of vegetable crops and its regulation. In: Yang C, Xu Y, Xie X, Wu Y, Gao Z, Li K et al, eds. *Growth Regulation and Quality Improvement of Vegetable Crops*. Singapore: Springer Nature Singapore; 2025. 495-557.

151. Al-Khayri JM, Rashmi R, Toppo V, Chole PB, Banadka A, Sudheer WN et al. Plant secondary metabolites: The weapons for biotic stress management. *Metabolites*. [Internet]. 2023;13(6). Available from: <https://doi.org/10.3390/metabo13060716>
152. Savatin DV, Gramegna G, Modesti V, Cervone F. Wounding in the plant tissue: The defense of a dangerous passage. *Front Plant Sci*. [Internet]. 2014;5. Available from: <https://doi.org/10.3389/fpls.2014.00470>
153. Farmer VC. Forest vegetation does recycle substantial amounts of silicon from and back to the soil solution with phytoliths as an intermediate phase, contrary to recent reports. *Eur J Soil Sci*. 2005;56(2):271-272.
154. Gfeller A, Baerenfaller K, Loscos J, Chételat A, Baginsky S, Farmer EE. Jasmonate controls polypeptide patterning in undamaged tissue in wounded arabidopsis leaves. *Plant Physiol*. 2011;156(4):1797-1807.
155. Liang YC, Ma TS, Li FJ, Feng YJ. Silicon availability and response of rice and wheat to silicon in calcareous soils. *Commun Soil Sci Plant Anal*. 1994;25:2285-2297.
156. Mitani Ueno N, Ogai H, Yamaji N, Ma JF. Physiological and molecular characterization of si uptake in wild rice species. *Physiol Plant*. 2013;151(3):200-207.
157. Gong H, Zhu X, Chen K, Wang S, Zhang C. Silicon alleviates oxidative damage of wheat plants in pots under drought. *Plant Sci*. 2005;169(2):313-321.
158. Liang Y, Wong J, Wei L. Silicon-mediated enhancement of cadmium tolerance in maize (*Zea mays* L.) grown in cadmium contaminated soil. *Chemosphere*. 2005;58(4):475-483.
159. Kim YH, Khan AL, Lee IJ. Silicon: A duo synergy for regulating crop growth and hormonal signaling under abiotic stress conditions. *Crit Revs Biotechnol*. 2015;36(6):1099-1109.
160. Shen S, Zhan C, Yang C, Fernie AR, Luo J. Metabolomics-centered mining of plant metabolic diversity and function: Past decade and future perspectives. *Mol Plant*. 2023;16(1):43-63.
161. Ma JF, Takahashi E. Soil, fertilizer and plant silicon research in Japan. Amsterdam, *The Netherlands: Elsevier Science*; 2002.
162. Ma JF, Yamaji N. Functions and transport of silicon in plants. *Cellular Mol Life Sci*. 2008;65:3049-3057.
163. Mir SH, Rashid I, Hussain B, Reshi ZA, Assad R, Sofi IA. Silicon supplementation of rescuegrass reduces herbivory by a grasshopper. *Front Plant Sci*. [Internet]. 2019;10. Available from: <https://doi.org/10.3389/fpls.2019.00671>
164. Bilal S, Shahzad R, Imran M, Jan R, Kim KM, Lee IJ. Synergistic association of endophytic fungi enhances *Glycine max* L. resilience to combined abiotic stresses: Heavy metals, high temperature and drought stress. *Ind Crops Prod*. [Internet]. 2020;143. Available from: <https://doi.org/10.1016/j.indcrop.2019.111931>
165. Zhou R, Yu X, Li X, Santos TMD, Rosenqvist E, Ottosen CO. Combined high light and heat stress induced complex response in tomato with better leaf cooling after heat priming. *Plant Physiol Biochem*. 2020;151:1-9.
166. Li Y, Cheng R, Spokas KA, Palmer AA, Borevitz JO. Genetic variation for life history sensitivity to seasonal warming in *Arabidopsis thaliana*. *Genetics*. 2014;196(2):569-577.
167. Lobell DB, Field CB. California perennial crops in a changing climate. *Clim Change*. 2011;109:317-333.
168. El-Naby SKMA, Abdelkhalek A, Baiea MHM, Amin OA. Impact of spraying some chemical substances on protecting Valencia orange trees from heat stress injuries. *Plant Arch*. 2020;20:2265-2270.
169. Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A et al. Crop production under drought and heat stress: Plant responses and management options. *Front Plant Sci*. [Internet]. 2017;8. Available from: <https://doi.org/10.3389/fpls.2017.01147>
170. Bhargava FFA. Morphological analysis of quinoa germplasm grown under lowland desert conditions. *J Agron Crop Sci*. 2010;197(2):124-134.
171. Lesjak J, Calderini DF. Increased night temperature negatively affects grain yield, biomass and grain number in Chilean quinoa. *Front Plant Sci*. [Internet]. 2017;8. Available from: <https://doi.org/10.3389/fpls.2017.00352>
172. Ghani MA, Abbas MM, Ali B, Ziaf K, Azam M, Anjum R et al. Role of salicylic acid in heat stress tolerance in tri-genomic *Brassica napus* L. *Bioagro*. 2020;33(1):13-20.
173. Knight H, Knight MR. Abiotic stress signalling pathways: Specificity and cross-talk. *Trends Plant Sci*. 2001;6(6):262-267.
174. El-Gayar S, Negm A, Abdrabbo M. Greenhouse operation and management in Egypt. In: Negm AM, ed. *Conventional Water Resources and Agriculture in Egypt*. Springer Cham: Springer International Publishing; 2018. 489-560.
175. Bakhat HF, Bibi N, Zia Z, Abbas S, Hammad HM, Fahad S et al. Silicon mitigates biotic stresses in crop plants: A review. *Crop Prot*. 2018;104:21-34.
176. Cooke J, Leishman MR. Consistent alleviation of abiotic stress with silicon addition: A meta analysis. *Funct Ecol*. 2016;30:1340-1357.
177. Hashem FA, El-Morshedy RM, Younis TM, Abdrabbo MAA. Foliar silicate application improves the tolerance of celery grown under heat stress conditions. *Acta Agric Slov*. [Internet]. 2021;117(3). Available from: <https://doi.org/10.14720/aas.2021.117.3.1865>
178. Ge H, Zhang F. Growth-promoting ability of *Rhodopseudomonas palustris* G5 and its effect on induced resistance in cucumber against salt stress. *J Plant Growth Regul*. 2018;38:180-188.
179. Fan Y, Zhang Y, Hess F, Huang B, Chen Z. Nutrient balance and soil changes in plastic greenhouse vegetable production. *Nutr Cycling Agroecosyst*. 2020;117:77-92.

180. Shalaby TA, Abd-Alkarim E, El-Aidy F, Hamed ES, Sharaf-Eldin M, Taha N et al. Nano-selenium, silicon and H_2O_2 boost growth and productivity of cucumber under combined salinity and heat stress. *Ecotoxicol Environ Saf.* [Internet]. 2021;212. Available from: <https://doi.org/10.1016/j.ecoenv.2021.111962>

181. Basirat M, Mousavi SM. Effect of foliar application of silicon and salicylic acid on regulation of yield and nutritional responses of greenhouse cucumber under high temperature. *J Plant Growth Regul.* 2022;41:1978-1988.

182. Abd-Elrahman SH, Hashem FAE, Farag AA. Mitigation of heat stress effects on snake cucumber plants by silicon additions and rice straw soil cover. *Commun Soil Sci Plant Anal.* 2022;53(6):759-772.

183. Bhandari R, Neupane N, Adhikari DP. Climatic change and its impact on tomato (*Lycopersicum esculentum* L.) production in plain area of Nepal. *Environ Challenges.* [Internet]. 2021;4. Available from: <https://doi.org/10.1016/j.envc.2021.100129>.

184. EL-AIDY F, ABDALLA M, EL-SAWY M-, KADY SA-E, BAYOUMI Y, EL-RAMADY H. Role of plant probiotics, sucrose and silicon in the production of tomato (*Solanum lycopersicum* L.) seedlings under heat stress in a greenhouse. *Appl Ecol Environ Res.* 2020;18(6):7685-7701.

185. Zakher AG, Abdrabbo MAA. Reduce the harmful effect of high temperature to improve the productivity of tomato under conditions of newly reclaimed land. *Egypt J Hortic.* 2014;41(2):151-167.

186. Soundararajan P, Sivanesan I, Jana S, Jeong BR. Influence of silicon supplementation on the growth and tolerance to high temperature in *Salvia splendens*. *Hortic, Environ, Biotechnol.* 2014;55:271-279.

187. El-Juhany LI. Degradation of date palm trees and date production in Arab countries: Causes and potential rehabilitation. *Aust J Basic Appl Sci.* 2010;4(8):3998-4010.

188. Patankar HV, Al-Harrasi I, Kharusi LA, Jana GA, Al-Yahyai R, Sunkar R et al. Overexpression of a Metallothionein 2A gene from date palm confers abiotic stress tolerance to yeast and *Arabidopsis thaliana*. *Int J Mol Sci.* [Internet]. 2019;20(12). Available from: <https://doi.org/10.3390/ijms20122871>

189. Khan A, Bilal S, Khan AL, Imran M, Shahzad R, Al-Harrasi A et al. Silicon and gibberellins: Synergistic function in harnessing ABA signaling and heat stress tolerance in date palm (*Phoenix dactylifera* L.). *Plants.* [Internet]. 2020;9(5). Available from: <https://doi.org/10.3390/plants9050620>

190. Mohapatra C, Chand R, Tiwari JK, Singh AK. Effect of heat stress during flowering and pod formation in pea (*Pisum sativum* L.). *Physiol Mol Biol Plants.* 2020;26:1119-1125.

191. Etesami H. Can interaction between silicon and plant growth promoting rhizobacteria benefit in alleviating abiotic and biotic stresses in crop plants? *Agric, Ecosyst Environ.* 2018;253:98-112.

192. Xu P, Guo Y, Bai J, Shang L, Wang X. Effects of long term chilling on ultrastructure and antioxidant activity in leaves of two cucumber cultivars under low light. *Physiol Plant.* 2008;132(4):467-478.

193. Kumar S, Malik J, Thakur P, Kaistha S, Sharma KD, Upadhyaya HD et al. Growth and metabolic responses of contrasting chickpea (*Cicer arietinum* L.) genotypes to chilling stress at reproductive phase. *Acta Physiol Plant.* 2010;33:779-787.

194. Molassiotis A, Sotiropoulos T, Tanou G, Diamantidis G, Therios I. Boron-induced oxidative damage and antioxidant and nucleolytic responses in shoot tips culture of the apple rootstock EM 9 (*Malus domestica* Borkh). *Environ Exp Bot.* 2006;56(1):54-62.

195. Bai X, Huang H, Li D, Yang F, Cong X, Wu S et al. Physiological and transcriptomic analyses reveal the role of the antioxidant system and jasmonic acid (JA) signal transduction in mulberry (*Morus alba* L.) response to flooding stress. *Horticulturae.* [Internet]. 2024;10(10). Available from: <https://doi.org/10.3390/horticulturae10101100>

196. Liang Y, Zhu J, Li Z, Chu G, Ding Y, Zhang J et al. Role of silicon in enhancing resistance to freezing stress in two contrasting winter wheat cultivars. *Environ Exp Bot.* 2008;64(3):286-294.

197. Waraich EA, Ahmad R, Ashraf MY, Saifullah, Ahmad M. Improving agricultural water use efficiency by nutrient management in crop plants. *Acta Agric Scand, Sect B — Soil Plant Sci.* 2011;61(4):291-304.

198. Habibi G. Role of trace elements in alleviating environmental stress. In: Ahmad P, Rasool S, eds. *Emerging Technologies and Management of Crop Stress Tolerance*. San Diego, CA, USA: Academic Press; 2014. 313-342.

199. Gunes A, Inal A, Bagci EG, Pilbeam DJ. Silicon-mediated changes of some physiological and enzymatic parameters symptomatic for oxidative stress in spinach and tomato grown in sodic-B toxic soil. *Plant Soil.* 2006;290:103-114.

200. Iwasaki K, Maier P, Fecht M, Horst WJ. Leaf apoplastic silicon enhances manganese tolerance of cowpea (*Vigna unguiculata*). *J Plant Physiol.* 2002;159(2):167-173.

201. Hsu YT, Kao CH. Cadmium-induced oxidative damage in rice leaves is reduced by polyamines. *Plant Soil.* 2007;291:27-37.

202. Wang LJ, Huang WD, Li JY, Liu YF, Shi YL. Peroxidation of membrane lipid and Ca^{2+} homeostasis in grape mesophyll cells during the process of cross-adaptation to temperature stresses. *Plant Sci.* 2004;167(1):71-77.

203. Hajiboland R. Silicon-mediated cold stress tolerance in plants. In: Saeedi HEHA, El-Ramady H, Fujita M, Pessarakli M, Hossain MA, eds. *Silicon and Nano-silicon in Environmental Stress Management and Crop Quality Improvement*. Oxford, United Kingdom: Academic Press; 2022. 161-180.

204. Epstein E. Silicon. *Annu Rev Plant Biol.* 1999;50:641-664.

205. Gunes A, Pilbeam DJ, Inal A, Bagci EG, Coban S. Influence of silicon on antioxidant mechanisms and lipid peroxidation in chickpea (*Cicer arietinum* L.) cultivars under drought stress. *J Plant Interact.* 2007;2(2):105-113.

206. Datnoff LE, Snyder GH, Korndörfer GH. *Silicon in Agriculture*. 1st ed. United Kingdom: Elsevier Science; 2001.

207. Fennell A. Freezing tolerance and injury in grapevines. *J Crop Improv.* 2004;10:201-235.

208. Habibi G. Effects of soil- and foliar-applied silicon on the resistance of grapevine plants to freezing stress. *Acta Biol Szegediensis.* 2015;50(2):109-117.

209. Habibi G. Exogenous silicon leads to increased antioxidant capacity in freezing-stressed pistachio leaves. *Acta Agric Slov.* 2015;105(1):43-52.

210. Luan H, Niu C, Nie X, Li Y, Wei M. Transcriptome and physiological analysis of rootstock types and silicon affecting cold tolerance of cucumber seedlings. *Plants.* [Internet]. 2022;11(3). Available from: <https://doi.org/10.3390/plants11030445>

211. Saad MM. Effect of some postharvest treatments on reducing chilling injury of cucumber fruits during cold storage. *Ann Agric Sci, Moshtohor.* 2019;57(2):455-468.

212. Song H, Yuan W, Jin P, Wang W, Wang X, Yang L et al. Effects of chitosan/nano-silica on postharvest quality and antioxidant capacity of loquat fruit during cold storage. *Postharvest Biol Technol.* 2016;119:41-48.

213. Shi S, Wang W, Liu L, Wu S, Wei Y, Li W. Effect of chitosan/nano-silica coating on the physicochemical characteristics of longan fruit under ambient temperature. *J Food Eng.* 2013;118(1):125-131.

214. Ghassemi S, Delangiz N, Lajayer BA, Saghafi D, Maggi F. Review and future prospects on the mechanisms related to cold stress resistance and tolerance in medicinal plants. *Acta Ecol Sin.* 2021;41(2):120-129.

215. Yin L, Wang S, Li J, Tanaka K, Oka M. Application of silicon improves salt tolerance through ameliorating osmotic and ionic stresses in the seedling of Sorghum bicolor. *Acta Physiol Plant.* 2013;35:3099-3107.

216. Sachdev S, Ansari SA, Ansari MI, Fujita M, Hasanuzzaman M. Abiotic stress and reactive oxygen species: Generation, signaling and defense mechanisms. *Antioxidants.* [Internet]. 2021;10(2). Available from: <https://doi.org/10.3390/antiox10020277>

217. Fujita M, Hasanuzzaman M. Approaches to enhancing antioxidant defense in plants. *Antioxidants.* [Internet]. 2022;11(5). Available from: <https://doi.org/10.3390/antiox11050925>

218. Zhu Z, Wei G, Li J, Qian Q, Yu J. Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.). *Plant Sci.* 2004;167(3):527-533.

219. Dhiman P, Rajora N, Bhardwaj S, Sudhakaran SS, Kumar A, Raturi G et al. Fascinating role of silicon to combat salinity stress in plants: An updated overview. *Plant Physiol Biochem.* 2021;162:110-123.

220. Rios JJ, Martínez-Ballesta MC, Ruiz JM, Blasco B, Carvajal M. Silicon-mediated improvement in plant salinity tolerance: The role of aquaporins. *Front Plant Sci.* [Internet]. 2017;8. Available from: <https://doi.org/10.3389/fpls.2017.00948>

221. Zhang W, Yu X, Li M, Lang D, Zhang X, Xie Z. Silicon promotes growth and root yield of *Glycyrrhiza uralensis* under salt and drought stresses through enhancing osmotic adjustment and regulating antioxidant metabolism. *Crop Prot.* 2018;107:1-11.

222. Pandey C, Khan E, Panthri M, Tripathi RD, Gupta M. Impact of silicon on Indian mustard (*Brassica juncea* L.) root traits by regulating growth parameters, cellular antioxidants and stress modulators under arsenic stress. *Plant Physiol Biochem.* 2016;104:216-225.

223. Laane HM. The effects of foliar sprays with different silicon compounds. *Plants.* [Internet]. 2018;7(2). Available from: <https://doi.org/10.3390/plants7020045>

224. Kovács S, Kutasy E, Csajbók J. The multiple role of silicon nutrition in alleviating environmental stresses in sustainable crop production. *Plants.* [Internet]. 2022;11(9). Available from: <https://doi.org/10.3390/plants11091223>